



# Aquaculturell

Prof. Dr. Hasan Hüseyin ATAR

- Extensive systems use low stocking densities (e.g., 5 000-10 000 shrimp post larvae (PL)/ha/crop) and no supplemental feeding, although fertilization may be done to stimulate the growth and production of natural food in the water. Water change is effected through tidal means, i.e., new water is let in only during high tide and the pond can be drained only at low tide. The ponds used for extensive culture are usually large (more than two ha) and may be shallow and not fully cleared of tree stumps. Production is generally low at less than 1 t/ha/y.



- Semi-intensive systems use densities higher than extensive systems (e.g., 50 000-100 000 shrimp PL/ha/crop) and use supplementary feeding. Intensive culture uses very high densities of culture organism (e.g., 200 000-300 000 shrimp PL/ha/crop) and is totally dependent on artificial, formulated feeds. Both systems use small pond compartments of up to one ha in size for ease of management.

- Semi-intensive and intensive culture systems are managed by the application of inputs (mainly feeds, fertilizers, lime, and pesticides) and the manipulation of the environment primarily by way of water management through the use of pumps and aerators. Feeding of the stock is done at regular intervals during the day.

- In intensive shrimp culture, the computed daily feed ration is given in equal doses from as low as three to as high as six times a day. Water change is also effected on a daily basis, with approximately 10-15% of the water in the pond replenished by the entry of new water in semi-intensive shrimp ponds.



- Semi-intensive and intensive culture systems are therefore more labour-intensive than extensive systems which need little attention, and are costlier to set up and operate, not to mention the fact that they also carry higher risks of mortalities resulting from disease, poor management, and/or force majeure (e.g., from anoxia due to non-functioning aerators during times of power failure).

- Production is of course much higher (for example, ranging from a minimum of 1.5 t/ha/crop from semi-intensive shrimp culture to a high of 10 t/ha/crop from intensive shrimp culture). Financial returns are therefore much more attractive than those from extensive culture, although studies have shown that the return on investment (ROI) from semi-intensive culture is better than from intensive culture due to the high cost of inputs (largely fry and feeds) used in intensive culture.



Parameter	Extensive	Semi-Intensive	Intensive
Species Used	Monoculture or Polyculture	Monoculture	Monoculture
Stocking Rate	Moderate	Higher than extensive culture	Maximum
Engineering Design and Layout	May or may not be well laid-out	With provisions for effective water management	Very well engineered system with pumps and aerators to control water quality and quantity
	Very big ponds	Manageable-sized units (up to 2 ha each)	Small ponds, usually 0.5-1 ha each
	Ponds may or may not be fully cleaned	Fully cleaned ponds	Fully cleaned ponds
Fertilizer	Used to enhance natural productivity	Used regularly with lime	Not used
Pesticides	Not used	Used regularly for prophylaxis	Used regularly for prophylaxis
Food and Feeding Regimen	None	Regular feeding of high quality feeds	Full feeding of high-quality feeds
		Depending on stocking density used, formulated feeds may be used partially or totally	
Cropping Frequency (crops/y)	2	2.5	2.5
Quality of Product	Good quality	Good quality	Good quality
	Culture species dominant but extraneous species may occur	Confined to culture species	Confined to culture species
	Variable sizes	Uniform sizes	Uniform sizes