

PEN303
QUANTUM MECHANICS

Prof. Dr. Handan OLĞAR

Course notes are prepared from the book Quantum Mechanics-Concepts and Applications by Zettili.

MATHEMATICAL TOOLS IN QUANTUM MECHANICS

- Hilbert Space and Wave Functions
- Dirac Notation
- Operators – General Definition
- Hermitian Adjoint
- Hermitian Operators
- Commutator Algebra

Hilbert Space and wave functions

A linear vector space consists of two sets of elements and two algebraic rules:

- a set of vectors ψ, ϕ, χ, \dots and a set of scalars a, b, c, \dots
- a rule for vector addition and a rule for scalar multiplication.
- Addition rule: The addition rule has the properties and structure of an abelian group
- Multiplication rule: The product of a scalar with a vector gives another vector.

Dirac Notation

- The physical state of a system is represented in quantum mechanics by elements of a Hilbert space; these elements are called state vectors. We can represent the state vectors in different bases by means of function expansions.
- To free state vectors from coordinate meaning, Dirac introduced what was to become an invaluable notation in quantum mechanics; it allows one to manipulate the formalism of quantum mechanics with ease and clarity.

- **Kets: elements of a vector space:** Dirac denoted the state vector ψ by the symbol $|\psi\rangle$, which he called a ket vector, or simply a ket. Kets belong to the Hilbert (vector) space H , or, in short, to the ket-space.
- **Bras: elements of a dual space:** Dirac denoted the elements of a dual space by the symbol $\langle|$, which he called a bra vector, or simply a bra; for instance, the element $\langle\psi|$ represents a bra.
- **Bra-ket:** Dirac notation for the scalar product:
- Dirac denoted the scalar (inner) product by the symbol $\langle| \rangle$, which he called a bra-ket. For instance, the scalar product ψ and ϕ is denoted by the bra-ket $\langle\psi|\phi\rangle$.

Operators

- General definition: An operator A is a mathematical rule that when applied to a ket $|\psi\rangle$ transforms it into another ket $|\psi'\rangle$ of the same space and when it acts on a bra $\langle\phi|$ transforms it into another bra : $\langle\phi'|$

$$\hat{A} |\psi\rangle = |\psi'\rangle, \quad \langle\phi| \hat{A} = \langle\phi'|.$$

Hermitian Adjoint

The Hermitian adjoint or conjugate², α^\dagger , of a complex number α is the complex conjugate of this number: $\alpha^\dagger = \alpha^*$. The Hermitian adjoint, or simply the adjoint, \hat{A}^\dagger , of an operator \hat{A} is defined by this relation:

$$\langle \psi | \hat{A}^\dagger | \phi \rangle = \langle \phi | \hat{A} | \psi \rangle^*.$$

Following these rules, we can write

$$(\hat{A}^\dagger)^\dagger = \hat{A},$$

$$(a\hat{A})^\dagger = a^* \hat{A}^\dagger,$$

$$(\hat{A}^n)^\dagger = (\hat{A}^\dagger)^n,$$

$$(\hat{A} + \hat{B} + \hat{C} + \hat{D})^\dagger = \hat{A}^\dagger + \hat{B}^\dagger + \hat{C}^\dagger + \hat{D}^\dagger,$$

$$(\hat{A}\hat{B}\hat{C}\hat{D})^\dagger = \hat{D}^\dagger\hat{C}^\dagger\hat{B}^\dagger\hat{A}^\dagger,$$

$$(\hat{A}\hat{B}\hat{C}\hat{D} | \psi)^\dagger = \langle \psi | \hat{D}^\dagger\hat{C}^\dagger\hat{B}^\dagger\hat{A}^\dagger.$$

- Further discussions about the mathematical concepts of Quantum Mechanics are discussed from the book Quantum Mechanics- Concepts and Applications by Zettili.