



A. Classifications of Dosage Forms Based on Route/ Method of Administration

The route of administration for a drug product is usually determined by the physicochemical characteristics of the drug molecule. Dosage forms are classified based on the route of administration into:

- a. Oral
- b. Topical
- c. Rectal
- d. Parenteral
- e. Respiratory/Inhaled
- f. Vaginal
- g. Ophthalmic
- h. Nasal
- i. Otic

a. Oral dosage forms

Oral dosage forms comprise pharmaceutical formulations taken orally for systemic effects. They are absorbed through the various epithelia and mucosa of the gastrointestinal tract at varying rates with the exception of drugs that are absorbed in the buccal cavity. Examples include tablets, capsules, suspensions, lozenges, pills, granules, powders, emulsions etc.



Photo credit: www.netdoctor.co.uk

b. Topical dosage forms

These include drug molecules that are in a suitable solid base (e.g., powders and aerosols), semi-solid base (e.g., ointments, creams, foams, gels, poultice and pastes), or in liquid form (e.g., solutions, suspension of solids in aqueous solutions or emulsions) which possesses either hydrophobic or hydrophilic properties. These drugs are applied to the skin or other topical surfaces (such as the eye, ear and nose) mainly for local action. Systemic drug delivery can also be achieved using topical preparations (e.g., transdermal patches), though absorption is often poor and erratic.



c. Rectal dosage forms

These are solutions, suppositories or emulsions administered rectally for local rather than systemic effect. These formulations can also be used to deliver drugs that are inactivated by gastrointestinal fluids when administered orally or when the oral route of the patient is precluded.



d. Parenteral dosage forms

These are usually sterile, particulate free and non-pyrogenic solutions or suspensions (of drugs in water or other suitable physiological acceptable vehicles) that are injected into the body using syringe and needle, infusion set etc.



e. Respiratory/Inhaled Dosage Forms

This is a type of dosage form where drugs are delivered in gaseous, aerosol mist or ultrafine solid particle form into the lungs. These classes of dosage form are mainly for direct treatment and management of respiratory diseases. Examples include nebulizers, powder aerosols and pressurized metered dose aerosols.



f. Vaginal dosage forms

These are dosage forms that are intended to be used in the vaginal cavity for either contraception, induction of labour, treatment of vaginal infections or local menopausal symptoms. Commonly used vaginal dosage forms include creams, tablets, vaginal gels and pessaries, suppositories, foams, ointments, tampons and inserts. Others include vaginal rings, vaginal films etc.



g. Ophthalmic Dosage Forms

These are principally sterile solutions, ointments and suspensions, essentially free from particles or substances that might irritate the eye. They are meant to be gently applied to the eyelids or placed in the pocket between the eyelids and the eyeball. Ophthalmic dosage forms are commonly used to treat local ocular disorders, e.g. infection and inflammation; or intraocular disorders e.g. glaucoma.



h. Nasal dosage forms

Nasal formulations are non-sterile aqueous-based systems that are instilled within or sprayed into the nasal cavity from a dropper or from a plastic squeeze bottle. They are predominantly employed for the treatment of local disorders – infections, congestion, and allergic rhinitis. Nasal preparations when absorbed through the nasal mucosa to achieve systemic effect.



i. Otic dosage forms

These are non-sterile aqueous solutions, or solutions prepared with glycerin or other solvents and dispersing agents that are instilled into the ear canal for the treatment of local disorders.





B. Classifications of Dosage Forms Based On the Physical Form of the Dosage Form

This class comprises the following dosage forms.

- a. Solid
- b. Semi-solid
- c. Liquid,
- d. Gaseous

a. Solid Dosage Forms

These comprise drug products with definite shape and volume. They constitute approximately 90% of all dosage forms clinically used to provide systemic administration of therapeutic agents. This class broadly encompasses two types of formulation – tablets and capsules. Others include powders, granules etc.



Photo credit: Shutterstock

b. Semi-solid dosage forms

These preparations applied on the skin or to the mucous membrane to achieve local or systemic effect. Examples include ointments, pastes, creams, gels etc. Semi-solid dosage forms have many characteristics in common – consistency, presentation, preservation requirement, and also route of administration which is mainly topical.



Photo Credit: www.e-konomista.pt

c. Liquid Dosage Forms

These include drug products administered in the form of solutions, suspensions, colloids, emulsions etc. Liquid dosage form can be sterile or non-sterile depending on the route of administration.



d. Gaseous dosage forms

This class comprises drug products that are packaged under pressure in a holder with a ceaseless or restricted conveyance valve framework. The gas inside contains restoratively dynamic medicaments that are released upon activation of an appropriate valve system. Examples include aerosols, nebulizer, sprays, inhalers etc.



General Considerations in Dosage Form Design

A suitable dosage form design includes

a. Preformulation Studies

These studies are designed to identify those physical and chemical properties of a candidate drug molecule which may affect the development of a safe, stable and efficient dosage forms with good bioavailability.

Commonly evaluated parameters during preformulation studies include – particle size and size distribution, solubility, dissolution behaviour, stability, refractive index, partition coefficient, drug-excipient compatibility, crystal form, surface properties, etc.



b. Biopharmaceutical Considerations

These studies are carried out to evaluate the rate and extent at which candidate drug molecule becomes available at the site of action. The aim is to achieve optimal therapeutic activity for the patient by modifying the delivery pattern of a drug molecule to systemic circulation.



The major biopharmaceutical considerations include

i. Pharmacodynamic Considerations

- Therapeutic objective.
- Toxic effect.
- Adverse reactions of candidate drug molecule.

ii. Drug Consideration

- Physicochemical characterization of the candidate drug molecules.

iii. Drug Product Consideration

- Bioavailability of candidate drug molecule.
- Pharmacokinetics of candidate drug molecule.
- Route of administration for the candidate drug molecule.
- Desired drug dosage form and
- Desired dose of the candidate drug molecule.

iv. Patient Consideration

- Compliance and acceptability of the final drug product

v. Manufacturing Considerations

- Cost
- Availability of pharmaceutical raw materials
- Stability and quality



c. Formulation and Development

This stage involves the actual combination of candidate drug molecule with various excipients and also optimizing the concentration at which each excipient is used. The choice of excipients depends on the properties of the drug molecule and the nature of the intended drug product.

