




NECROSIS



NECROSIS VS APOPTOSIS

- **Necrosis** is the death of cells or organs within living organism.
- In contrast, **apoptosis** is a naturally occurring **programmed** and targeted cause of cellular death(see slides 146-150).
- While **apoptosis** often provides **beneficial** effects to the organism, necrosis is almost always **detrimental** and can be **fatal**



Pathologic Basis of Veterinary Disease, 6th Ed., Ed.: Zachary JF, (2017) Elsevier, St. Louis, Missouri.


NECROSIS

Causes of necrosis

- Necrosis is caused by :
 - ✓ **Chemical substances** such as mineral acids, caustic alkalines and phenol.
 - ✓ **Physical influences** such as heat, cold, electricity
 - ✓ **Mechanical forces**
 - ✓ **Nutrition** (failure of blood to reach the organs because of thrombosis, embolism, volvulus, invagination, tumor, abscess, etc.)
 - ✓ **Neural causes** (peripheral nerve destruction)
 - ✓ **Infectious causes** (bacterial: necrobacillosis, tuberculosis, pseudotuberculosis. viral: rinderpest, foot and mouth disease. mycotic: aspergillosis, blastomycosis)

NUCLEAR CHANGES IN NECROSIS

- Four types of nuclear changes may occur:
- ✓ **Pyknosis** : condensation of nuclear chromatin into a dark round, homogeneous mass smaller than in a normal nucleus.
- ✓ **Karyorrhexis** : breaking up of the nucleus into numerous pieces.
- ✓ **Karyolysis** : dissolution of nuclear chromatin, leaving a ghost form of the nucleus
- ✓ And finally : **complete absence of the nucleus.**



A




Figure 1-16 Cytoarchitecture of Cellular Necrosis. A, Schematic representation of nuclear and cytoplasmic changes in the stages of necrosis. rER, rough endoplasmic reticulum. B, Pyknosis and karyolysis, renal tubular epithelium toxicosis, mouse. Some tubular epithelial cells have undergone hydropic degeneration; others are necrotic with pyknosis (arrow) or karyolysis (arrowhead). H&E stain. C, Karyorrhexis, lymphocytes, spleen, dog. Necrotic lymphocytes and free nuclei (arrow) because of parvovirus infection. H&E stain. (A courtesy Dr. M.A. Miller, College of Veterinary Medicine, Purdue University; and Dr. J.F. Zachary, College of Veterinary Medicine, University of Illinois; B and C courtesy Dr. L.H. Arp.)

TYPES OF NECROSIS

- Coagulative necrosis
 - Caseous necrosis
- Liquefactive necrosis
- Gangrene


COAGULATIVE NECROSIS

- The **denaturation** of cytoplasmic **proteins**, which at the histologic level imparts an opaque and **intense cytoplasmic eosinophilia** to necrotic cells.
- Coagulative necrosis is a typical early response to *hypoxia, ischemia, or toxic injury*.
- Occurs in **infarcts** of the kidney, spleen and myocardium.
- **Grossly**, the area of necrotic tissue is yellowish grey or pale yellow in color and is depressed below the surface of the surrounding tissue.
- **Microscopically**, the architectural outline of the tissue is preserved but the cellular detail is lost.

CASEOUS NECROSIS


- **Caseous**, from the Latin word for **cheese**, refers to the curdled or cheese like gross appearance of this form of necrosis.
- Characterized by the **loss of both architectural and cellular detail**.
- **Macroscopically**, caseation may appear as crumbled, granular, or laminated yellow-white exudate in the center of a granuloma or a chronic abscess.
- **Histologically**, the lysis of leukocytes and parenchymal cells converts the necrotic tissue into a granular to amorphous eosinophilic substance with basophilic nuclear debris.
- This type of necrosis is characteristic of the lesions of **tuberculosis**.

CASEOUS NECROSIS




Tuberculosis


CASEOUS NECROSIS




wedge shape infarct in kidney (coagulative necrosis)



www.medicosite.com



www.medicosite.com



LIQUEFACTIVE NECROSIS

- Liquefactive necrosis is typical of organs **in which the tissues have a lot of lipid**.
- Its often seen **in brain** and **spinal cord** and the term for the macroscopic (gross) appearance of necrosis in the brain and spinal cord is **malacia**.
- In liquefactive necrosis, cells are lysed, and the necrotic tissue is converted to a fluid phase.
- Grossly, the dead tissue is in a liquid form and sometimes *has a creamy or pasty consistency*.
- Microscopically, *empty spaces are seen since the liquefied material are lost during the processing of the tissue*. A pinkish staining protein material may remain or may not as a precipitate from the liquid.

GANGRENE

- Gangrene denotes a type of necrosis that tends to develop at the distal aspect of extremities, such as the limbs, tail, or pinnae, or in dependent portions of organs, such as the mammary glands or lung lobes.
- Gangrene can be designated as **wet** or **dry**.
- **Wet gangrene** occurs if the necrotic tissue is infected by *certain bacteria*.
- ✓ Hydrogen sulfide is produced
- ✓ As a result of hydrogen sulfide binding with blood pigments, the tissues become greenish-black in color.
- ✓ E.g. Septic ulcers of the uterus, gangrenous pneumonia

GANGRENE

- If those bacteria are *gas forming* (e.g., *Clostridium* spp.), then wet gangrene becomes **gas gangrene**.
 - ✓ E.g. Blackleg and malignant edema.
- **Dry gangrene** is the result of decreased vascular perfusion and/or loss of blood supply.
 - ✓ It is a form of infarction resulting in coagulative necrosis that imparts a dry, leathery texture to the necrotic tissue, providing that it remains **free of putrefactive bacteria**.
 - ✓ Necrosis in case of tearing of the umbilical cord after birth.
 - ✓ Dry dying fetus in utero.

RESULTS OF NECROSIS

- **Dissolution** and **resorption** of necrotic tissue.
- Abscess formation
- Sloughing and desquamation of the necrotic tissue
- **Encapsulation** without liquefaction ; a containing wall of leukocytes and connective tissue is formed around the mass and the area become encapsulated.
- Organisation of necrotic tissue: occurs when necrotic tissue is invaded with capillaries, connective tissue and leucocytes. The necrotic tissue is digested by leukocytes then the areas are infiltrated **with connective tissue** leaving **a scar**.
- **Calcification** of necrotic material
- **Gangrene**
- **Atrophy**
- **Regeneration**

INFARCTUS

Infarction is a local area of ischemic necrosis in a tissue caused by occlusion of the arterial supply or venous drainage.


It is usually formed in organs with insufficient veins, without collateral connections.

As a result of embolism, thrombosis or spasm, a complete blockage of the artery does not lead to the region where the endothelium feeds, and ischemia (local anemia) develops.

The coagulation necrosis on the side of the occluded vein is called infarct.

In the formation of infarctus

- The organ is resistant to oxygen (for example, the kidneys are unstable, the infarctus is seen)
- Whether blood circulation, vascularisation is sufficient,
- collateral vascularisations are not adequately fed and fed on other vessels (eg infarctus does not occur due to excessive vascular anastomoses in skeletal muscles, skin, uterus, thyroid tissue).



ACCORDING TO MORPHOLOGY

Anemic infarction

- Artery blockage in organs such as the heart, kidney, spleen, brain is the result. It appears pale from the beginning.

Hemorrhagic infarction

- Occurs when the venous system is obstructed. This is most commonly seen in the lungs.
- It also occurs in the testes and ovaries. It is reddish due to blood stagnation

ACCORDING TO TIME

- New (acute), old (chronic)

MORPHOLOGICAL APPEARANCE

- Coagulation NECROSIS is occurred
- Lungs, heart-like organs do not have a specific shape. It's like a map and its boundaries are irregular.
- It is wedge-shaped (like triangular). in the kidneys and
- the spleen.
- Infarcts tend to be wedge-shaped; the base of the wedge is at periphery, with the occluded vessel at the apex. The margins of the infarct may be irregular, a reflection of the vascular supply adjacent, nonaffected tissue.
- However, almost six of the capsules are not significantly affected by being fed with blood vessels in the capsule.