



LENFOİD SİSTEME GİRİŞ VE LENFOİD ORGAN PATOLOJİLERİ

Prof. Dr. Işın Kuzu
Patoloji Anabilim Dalı
Dönem III Modül IV
13.03.2017




HEMATOPOIESIS



HEMATOPOIEN

KEMİK İLİĞİ

Robbins Cotran Kumar



ERİTROPOEZ

- Proeritroblast
- Erken eritroblast
- İntermedier eritroblast
- Geç eritroblast
- Retikülosit
- Eritrosit




GRANULOPOEZ


- Myeloblast
- Promyelosit
- Myelosit
- Metamyelosit
- Çomak
- Polimorf nüveli lökosit
 - Bazofil
 - Nötrofil
 - Eosinofil

MEGAKARYOPOEZ VE TROMBOPOEZ

- Megakaryoblast
- Promegakaryosit
- Megakaryosit
- Trombosit

KEMİK İLİĞİ HÜCRELERİNİN ORANI VE DAĞILIMI





HEMATOPOIEN

KANDAKI HÜCRELERİN ORANLARI

TABLE 13-1 Adult Reference Ranges for Blood Cells*

Cell Type	
White cells ($\times 10^3/\mu\text{L}$)	4.8-10.8
Granulocytes (%)	40-70
Neutrophils ($\times 10^3/\mu\text{L}$)	1.4-6.5
Lymphocytes ($\times 10^3/\mu\text{L}$)	1.2-3.4
Monocytes ($\times 10^3/\mu\text{L}$)	0.1-0.6
Eosinophils ($\times 10^3/\mu\text{L}$)	0-0.5
Basophils ($\times 10^3/\mu\text{L}$)	0-0.2
Red cells ($\times 10^3/\mu\text{L}$)	4.3-5, men; 3.5-5, women
Platelets ($\times 10^3/\mu\text{L}$)	150-450


TABLE 13-2 Mechanisms and Causes of Leukocytosis

Increased Production in the Marrow
Chronic infection or inflammation (growth factor-dependent) Paraneoplastic (e.g., Hodgkin lymphoma; growth factor-dependent) Myeloproliferative disorders (e.g., chronic myeloid leukemia; growth factor-independent)
Increased Release from Marrow Stores
Endotoxemia Infection Hypoxia
Decreased Margination
Exercise Catecholamines
Decreased Extravasation into Tissues
Glucocorticoids

LÖKOSİTOZ

Type of Leukocytosis	Causes
Neutrophilic leukocytosis	Acute bacterial infections, especially those caused by pyogenic organisms; sterile inflammation caused by, for example, tissue necrosis (myocardial infarction, burns)
Eosinophilic leukocytosis (eosinophilia)	Allergic disorders such as asthma, hay fever, parasitic infestations; drug reactions; certain malignancies (e.g., Hodgkin and some non-Hodgkin lymphomas); autoimmune disorders (e.g., pemphigus, dermatitis herpetiformis) and some vasculitides; atheroembolic disease (transient)
Basophilic leukocytosis (basophilia)	Rare, often indicative of a myeloproliferative disease (e.g., chronic myelogenous leukemia)
Monocytosis	Chronic infections (e.g., tuberculosis), bacterial endocarditis, rickettsiosis, and malaria; autoimmune disorders (e.g., systemic lupus erythematosus); inflammatory bowel diseases (e.g., ulcerative colitis)
Lymphocytosis	Accompanies monocytosis in many disorders associated with chronic immunologic stimulation (e.g., tuberculosis, brucellosis); viral infections (e.g., hepatitis A, cytomegalovirus, Epstein-Barr virus); <i>Bordetella pertussis</i> infection

LENFOSİT OLUŞUMU VE BAĞIŞIKLIK YANITI



LENFOSİT AİLESİ


Class	Functions	Antigen receptor and specificity	Selected markers	Percent of total lymphocytes (human)		
				Blood	Lymph node	Spleen
ab T lymphocytes						
CD4 ⁺ helper T lymphocytes	B cell differentiation (humoral immunity) Macrophage activation (cell-mediated immunity)	ab heterodimers Diverse specificities for peptide-class II MHC complexes	CD3 ⁺ , CD4 ⁺ , CD8 [−]	50–60*	50–60	50–60
CD8 ⁺ cytotoxic T lymphocytes	Killing of cell infected with microbes, killing of tumor cells	ab heterodimers Diverse specificities for peptide-class I MHC complexes	CD3 ⁺ , CD4 [−] , CD8 ⁺	20–25	15–20	10–15
Regulatory T cells	Suppress function of other T cells (regulation of immune responses, maintenance of self-tolerance)	ab heterodimers	CD3 ⁺ , CD4 ⁺ , CD25 ⁺ (Most common, but other phenotypes as well)	Rare	10	10
gd T lymphocytes	Helper and cytotoxic functions (innate immunity)	gd heterodimers Limited specificities for peptide and nonpeptide antigens	CD3 ⁺ , CD4, and CD8 variable			
B lymphocytes	Antibody production (humoral immunity)	Surface antibody Diverse specificities for all types of molecules	Fc receptors; class II MHC; CD19; CD21	10–15	20–25	40–45
Natural killer cells	Killing of virus-infected or damaged cells (innate immunity)	Various activating and inhibitory receptors Limited specificities for MHC or MHC-like molecules	CD16 (Fc receptor for IgG)	10	Rare	10
NKT cells	Suppress or activate innate and adaptive immune responses	ab heterodimers (Limited specificity for glycolipid-CD1 complexes)	CD16 (Fc receptor for IgG); CD3	10	Rare	10

*In most cases, the ratio of CD4⁺CD8⁻ to CD8⁺CD4⁻ cells is about 2:1.
Abbreviations: IgG, immunoglobulin G; MHC, major histocompatibility complex


NAİVE EFEKTÖR VE BELLEK LENFOSİT KARAKTERLERİ

	Naive lymphocytes	Activated or effector lymphocytes	Memory lymphocytes
T lymphocytes			
Migration	Preferentially to peripheral lymphoid tissues	Preferentially to inflamed tissues	Preferentially to inflamed tissues, mucosal tissues
Frequency of cells responsive to particular antigen	Very low	High	Low
Effector functions	None	Cytokine secretion; cytotoxic activity	None
Cell cycling	No	Yes	+/-
Surface protein expression			
High-affinity IL-2 receptor	Low	High	Low
Peripheral lymph node homing receptor (L-selectin, CD62L)	High	Low	Low or variable
Adhesion molecules: integrins, CD44	Low	High	High
Chemokine receptor: CCR7	High	Low	Variable
Major CD45 isoform (humans only)	CD45RA	CD45RO	CD45RO; variable
Morphology	Small; scant cytoplasm	Large; more cytoplasm	Small
B lymphocytes			
Membrane immunoglobulin (Ig) isotype	IgM and IgD	Frequently IgG, IgA, IgE	Frequently IgG, IgA, IgE
Affinity of Ig produced	Relatively low	Increases during immune response	Relatively high
Effector function	None	Antibody secretion	None
Morphology	Small; scant cytoplasm	Large; more cytoplasm; some are plasma cells	Small
Chemokine receptor: CXCR5	High	Low	?
CD27	Low	High	High

Abbreviations: IL-2, interleukin-2



© Elsevier. Abbas et al: Cellular and Molecular Immunology 6e - www.studentconsult.com




© Elsevier. Abbas et al: Cellular and Molecular Immunology 6e - www.studentconsult.com

LENFOİD DOKULAR

- LENF NODÜLLERİ VE LENFATİK SİSTEM
- DALAK
- DERİ
- MUKOZAL İMMUN SİSTEM



KEMİK İLİĞİ
TİMUS

LENF NODÜLLERİ
LENFATİK SİSTEM
DALAK
DERİ
MUKOZAL İMMUN SİSTEM




TİMUS




T LENFOSİT
OLGUNLAŞMA VE
İMMUN TOLERANS
GELİŞİMİ





LENF NODÜLLERİ VE LENFATİK SİSTEM




LENF NODÜLLERİ




LENF NODÜLLERİ




DERI




MUKOZAL IMMUN SISTEM




LENFOSİT DOLAŞIMI YOLLARI




HIGH ENDOTHELIAL VENÜLLER




(B) L-selectin ligand on endothelial cells




(C) T cells binding to HEV: frozen section assay



(D) T cells binding to HEV: electron micrograph



NAIVE VE EFEKTÖR T LENFOSİTLERİN GÖÇÜ







B

T cell homing receptor	Ligand on endothelial cell	Function of receptor: ligand pair
Naive T cells L-selectin CCR7	L-selectin ligand CCL19 or CCL21	Adhesion of naive T cells to high endothelial venule in lymph node
Activated (effector and memory) T cells E- and P-selectin ligand LFA-1 (β2-integrin) or VLA-4 (β1 integrin) CXCR3 CCR5	E- or P-selectin ICAM-1 or VCAM-1 CXCL10 (others) CCL4 (others)	Initial weak adhesion of effector and memory T cells to cytokine activated endothelium at peripheral site of infection Stable arrest on cytokine activated endothelium at peripheral site of infection Activation of integrins and chemokinesis Activation of integrins and chemokinesis

DALAK


Çevre kanının en önemli antikor kaynağı



DALAK İŞLEVLERİ

- *Kan hücre ve partiküllerin fagositozu*
- *Antikor yapımı*
- *Hematopoez*
- *Kan hücrelerinin yıkımı (sekestrasyon)*

DALAK



SPLENOMEGALI

TABLE 13-12 Disorders Associated with Splenomegaly

I. Infections

Nonspecific splenitis of various blood-borne infections (particularly infectious endocarditis)

Infectious mononucleosis

Tuberculosis

Typhoid fever

Brucellosis

Cytomegalovirus

Syphilis

Malaria

Histoplasmosis

Toxoplasmosis

Kala-azar

Trypanosomiasis

Schistosomiasis

Leishmaniasis

Echinococcosis

SPLENOMEGALI

II. Congestive States Related to Portal Hypertension

Cirrhosis of the liver
Portal or splenic vein thrombosis
Cardiac failure

III. Lymphohematogenous Disorders

Hodgkin lymphoma
Non-Hodgkin lymphomas and lymphocytic leukemias
Multiple myeloma
Myeloproliferative disorders
Hemolytic anemias

IV. Immunologic-Inflammatory Conditions

Rheumatoid arthritis
Systemic lupus erythematosus

V. Storage Diseases

Gaucher disease
Niemann-Pick disease
Mucopolysaccharidoses

VI. Miscellaneous Disorders

Amyloidosis
Primary neoplasms and cysts
Secondary neoplasms

LENFADENOPATİ

- 1- Reaktif veya
 infeksiyonlara bağlı
- 2- Neoplastik büyümeler
 (Primer - sekonder
 (metastatik))

Romatolojik - immünolojik

Romatoid artrit

Lupus eritematosus





Sjögren sendromu

İmmün yetmezlik durumları..


Enfeksiyöz

Bakterial


Viral




Granülomatöz inflamasyon




Granüloamatöz inflamasyon (nekroz)





Sarkoidozis




Sarkoidozis nedeni bilinmeyen ve birden fazla sistemi etkileyen, nonkazeöz granülomatöz hastalıktır

Akciğer infiltrasyonları, malignite ile karışabilen hiler lenfadenopati


Lenf düğümünde tüberküloz ile karışabilecek granülomatöz inflamasyon oluşturur.

Sarkoidosis



Süpüratif granülomatöz lenfadenopatiler:
düzensiz kenarlı nekroz alanları ile karakterli,
Tüberküloz ile karışabilecek morfolojiye sahip
bir grup hastalık

Atipik mikobakteri infeksiyonları,
Kedi tırmığı hastalığı (Bartonella henselae)
Tularemi,
Yersinia,
Lenfogranüloma venereum (Chlamidia
trachomatis)




LENFOTROP VİRÜSLER

Human Herpes Virüsler (ds DNA)



- EBV (HHV4) Enfeksiyöz Mononükleozis
- KSV (Kaposi Sarkoma Virus) (HHV8)
Castleman Hastalığı
- CMV (Sitomegalovirus) (HHV5)
- HIV (ds RNA)

Lenfoid Hücreler




Lenfoid Hücreler

Hümmoral Bağışıklık





B hücre
Reseptörü
İmmünglobulin
(Ig) Molekülü




Lenfoid Hücreler

Hümmoral Bağışıklık




B hücre
Reseptörü
İmmünglobulin
(Ig) Molekülü






Lenfoid Hücreler

Hümmoral Bağışıklık




B hücre
Reseptörü
İmmünglobulin
(Ig) Molekülü




T Hücre
reseptörü
TCR
Molekülü

Hüccresel immunit





Lenfoid Hücreler

Hümmoral Bağışıklık




Cedar, Nat Rev Immunol 2015
Björkström Nat Rev Immunol 2016
Abbas, Cellular & Molecular Immunology


**B hücre
Reseptörü
İmmünglobulin
(Ig) Molekülü**





**T Hücre
reseptörü
TCR
Molekülü**



Hüccresel Bağışıklık





**Doğal
Bağışıklık**



HEMATOLOJİK NEOPLAZİLERİN GELİŞME MEKANİZMALARI

- NORMAL HÜCRELERİN OLGUNLAŞMA, BÜYÜME, YAŞAMA VE ÖLÜM İŞLEVLERİYLE İLGİLİ MOLEKÜLLER (SİTOKİN – BÜYÜME FAKTÖRLERİ)
- HIZLI ÇOĞALMA GEREKLİLİĞİ
- KALITSAL FAKTÖRLER
- ENFEKSİYONLAR
- KRONİK İNFLAMASYONLAR
- KİMYASAL VE FİZİKSEL KARSİNOJENLER (ilaçlar, radyasyon, sigara, çeşitli diğer kimyasal karsinojenler)





Tümör gelişimi ile ilişkili onkogenler

Category	Proto-Oncogene	Mode of Activation in Tumor	Associated Human Tumor
Growth Factors			
PDGF-β chain	<i>PDGFB</i>	Overexpression	Astrocytoma
Fibroblast growth factors	<i>HST1</i>	Overexpression	Osteosarcoma
	<i>FGF3</i>	Amplification	Stomach cancer
			Bladder cancer
			Breast cancer
			Melanoma
TGF-α	<i>TGFA</i>	Overexpression	Astrocytomas
HGF	<i>HGF</i>	Overexpression	Hepatocellular carcinomas
			Thyroid cancer
Growth Factor Receptors			
EGF-receptor family	<i>ERBB1 (EGFR)</i>	Mutation Amplification	Adenocarcinoma of lung
	<i>ERBB2 (HER)</i>		Breast carcinoma
FMS-like tyrosine kinase 3	<i>FLT3</i>	Point mutation	Leukemia

Category	Proto-Oncogene	Mode of Activation in Tumor	Associated Human Tumor
Receptor for neurotrophic factors	<i>RET</i>	Point mutation	Multiple endocrine neoplasia 2A and B, familial medullary thyroid carcinomas
PDGF receptor	<i>PDGFRB</i>	Overexpression, translocation	Gliomas, leukemias
Receptor for KIT ligand	<i>KIT</i>	Point mutation	Gastrointestinal stromal tumors, seminomas, leukemias
ALK receptor	<i>ALK</i>	Translocation, fusion gene formation	Adenocarcinoma of lung, certain lymphomas
		Point mutation	Neuroblastoma
Proteins Involved in Signal Transduction			
GTP-binding (G) proteins	<i>KRAS</i>	Point mutation	Colon, lung, and pancreatic tumors
	<i>HRAS</i>	Point mutation	Bladder and kidney tumors
	<i>NRAS</i>	Point mutation	Melanomas, hematologic malignancies
	<i>GNAQ</i>	Point mutation	Uveal melanoma
	<i>GNAS</i>	Point mutation	Pituitary adenoma, other endocrine tumors
Nonreceptor tyrosine kinase	<i>ABL</i>	Translocation	Chronic myelogenous leukemia
		Point mutation	Acute lymphoblastic leukemia

Category	Proto-Oncogene	Mode of Activation in Tumor	Associated Human Tumor
RAS signal transduction	<i>BRAF</i>	Point mutation, Translocation	Melanomas, leukemias, colon carcinoma, others
Notch signal transduction	<i>NOTCH1</i>	Point mutation, Translocation Gene rearrangement	Leukemias, lymphomas, breast carcinoma
JAK/STAT signal transduction	<i>JAK2</i>	Translocation	Myeloproliferative disorders
			Acute lymphoblastic leukemia
Nuclear Regulatory Proteins			
Transcriptional activators	<i>MYC</i>	Translocation	Burkitt lymphoma
	<i>NMYC</i>	Amplification	Neuroblastoma
Cell Cycle Regulators			
Cyclins	<i>CCND1</i> (Cyclin D1)	Translocation	Mantle cell lymphoma, multiple myeloma
		Amplification	Breast and esophageal cancers
Cyclin-dependent kinase	<i>CDK4</i>	Amplification or point mutation	Glioblastoma, melanoma, sarcoma

HEMATOLOJİK NEOPLAZİLER

MİYELOİD NEOPLAZİLER

- MİYELODİSPLASTİK SENDROMLAR
- AKUT MİYELOİD LÖSEMİLER
- KRONİK MİYELOPROLİFERATİF NEOPLAZİLER

LENFOMALAR

- LENFOBLASTİK LÖSEMİLER (T ve B Hücreli)
- HODGKİN DIŞI LENFOMALAR :
 - 1- OLGUN B HÜCRELİ LENFOMALAR - PLAZMA HÜCRELİ NEOPLAZİLER
 - 2- OLGUN T HÜCRELİ LENFOMALAR
- HODGKİN LENFOMA

HİSTİYOSİTİK NEOPLAZİLER