MTH 102 Calculus -II

Dr. Gizem SEYHAN ÖZTEPE Topic-9-Series



Series

An infinite series (or series) is the sum of an infinite sequence of numbers

$$a_1 + a_2 + \cdots + a_n + \cdots$$

The goal of this section is to understand the meaning of such an infinite sum and to develop methods to calculate it. The number a_n is the nth term of the series. The sequence $\{s_n\}$ defined by

$$s_1 = a_1$$

$$s_2 = a_1 + a_2$$

...

$$s_n = a_1 + a_2 + \dots + a_n = \sum_{k=1}^n a_k$$

is the sequence of partial sums of the series, the number s_n being the nth partial sum.



If the sequence of partial sums converges to a limit L, we say that the series converges and that its sum is L. In this case we also write

$$a_1 + a_2 + \dots + a_n + \dots = \sum_{n=1}^{\infty} a_n = L$$

If the sequence of partial sums of the series does not converge, we say that the series diverges.

Thus the sum of a series is the limit of the sequence of partial sums. So when we write $\sum_{n=1}^{\infty} a_n = s$, we mean that by adding sufficiently many terms of the series we can get as close as we like to the number s. Notice that

$$\sum_{n=1}^{\infty} a_n = \lim_{n \to \infty} \sum_{i=1}^{n} a_i$$



4 The geometric series

$$\sum_{n=1}^{\infty} ar^{n-1} = a + ar + ar^2 + \cdots$$

is convergent if |r| < 1 and its sum is

$$\sum_{n=1}^{\infty} ar^{n-1} = \frac{a}{1-r} \qquad |r| < 1$$

If $|r| \ge 1$, the geometric series is divergent.

Each term is obtained from the preceding one by multiplying it by the common ratio .

In words: The sum of a convergent geometric series is



Some properties of the series are as follows:

8 Theorem If Σ a_n and Σ b_n are convergent series, then so are the series Σ ca_n (where c is a constant), Σ $(a_n + b_n)$, and Σ $(a_n - b_n)$, and

(i)
$$\sum_{n=1}^{\infty} ca_n = c \sum_{n=1}^{\infty} a_n$$

(ii)
$$\sum_{n=1}^{\infty} (a_n + b_n) = \sum_{n=1}^{\infty} a_n + \sum_{n=1}^{\infty} b_n$$

(iii)
$$\sum_{n=1}^{\infty} (a_n - b_n) = \sum_{n=1}^{\infty} a_n - \sum_{n=1}^{\infty} b_n$$

