


Physics 122: Electricity & Magnetism – Lecture 3 Coulomb's Law

Prof.Dr. Barış Akaoglu

Electric Force and Coulomb's Law

- We can measure the force of attraction or repulsion between charges, call them q_1 and q_2 (we will use the symbol q or Q for charge).



- When we do that, we find that the force is proportional to the each of the charges, is inversely proportional to the distance between them, and is directed along the line between them (along r).



- In symbols, the magnitude of the force is $F = k \frac{|q_1||q_2|}{r^2}$ where k is some constant of proportionality.
- This force law was first studied by Coulomb in 1785, and is called Coulomb's Law. The constant $k = 8.98755 \times 10^9 \text{ N m}^2/\text{C}^2$ is the Coulomb constant.

Electric Force and Coulomb's Law

- Although we can write down a vector form for the force, it is easier to simply use the equation for the magnitude, and just use the "like charges repel, opposites attract" rule to figure out the direction of the force.
- Note that the form for Coulomb's Law is exactly the same as for gravitational force between two masses

$$F = G \frac{m_1 m_2}{r^2}$$

$$G \Rightarrow k$$
$$m \Rightarrow q$$

Note BIG difference, There is only one "sign" of mass, only attraction.

- Note also that the mass is an intrinsic property of matter. Likewise, charge is also an intrinsic property. We only know it exists, and can learn its properties, because of the force it exerts.

Full form of Coulomb's Law

- Because it makes other equations easier to write, Coulomb's constant is actually written


$$k = \frac{1}{4\pi\epsilon_0}$$

$$F = \frac{1}{4\pi\epsilon_0} \frac{|q_1||q_2|}{r^2}$$

where $\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/\text{N}\cdot\text{m}^2$ is called the permittivity constant.

Spherical Conductors

- Because it is conducting, charge on a metal sphere will go everywhere over the surface.
- You can easily see why, because each of the charges pushes on the others so that they all move apart as far as they can go. Because of the symmetry of the situation, they spread themselves out uniformly.
- There is a theorem that applies to this case, called the shell theorem, that states that the sphere will act as if all of the charge were concentrated at the center.




Note, forces are equal and opposite



These two situations are the same

Case of Multiple Charges

- You can determine the force on a particular charge by adding up all of the forces from each charge.



Forces on one charge due to a number of other charges

Let's Calculate the Exact Location

- Force is attractive toward both negative charges, hence could balance.
- Need a coordinate system, so choose total distance as L , and position of $+$ charge from $-q$ charge as x .
- Force is sum of the two force vectors, and has to be zero, so


$$F = F_1 + F_2 = k \frac{2qQ}{(L-x)^2} - k \frac{qQ}{x^2} = 0$$

- A lot of things cancel, including Q , so our answer does not depend on knowing the $+$ charge value. We end up with

$$\frac{2}{(L-x)^2} = \frac{1}{x^2}$$

$$\frac{(L-x)^2}{x^2} = 2 \Rightarrow \frac{L-x}{x} = \sqrt{2}$$

- Solving for x , $x = \frac{L}{1+\sqrt{2}} = 0.412L$, so slightly less than half-way between.



Summary

- Charge is an intrinsic property of matter.
- Charge comes in two opposite senses, positive and negative.
- Mobile charges we will usually deal with are electrons, which can be removed from an atom to make positive charge, or added to an atom to make negative charge. A positively charged atom or molecule can also be mobile.
- There is a smallest unit of charge, e , which is $e = 1.602 \times 10^{-19}$ C. Charge can only come in units of e , so charge is *quantized*. The unit of charge is the Coulomb.
- Charge is *conserved*. Charge can be destroyed only in pairs ($+e$ and $-e$ can annihilate each other). Otherwise, it can only be moved from place to place.
- Like charges repel, opposite charges attract.
- The electric force is given by Coulomb's Law:
$$F = \frac{1}{4\pi\epsilon_0} \frac{|q_1||q_2|}{r^2}$$
- Materials can be either conductors or insulators.
- Conductors and insulators can both be charged by adding charge, but charge can also be *induced*.
- Spherical conductors act as if all of the charge on their surface were concentrated at their centers.