Overview: The Body's Long-Distance Regulators

- Animal hormones are chemical signals that are secreted into the circulatory system and communicate regulatory messages within the body.
- Hormones reach all parts of the body, but only target cells are equipped to respond.
- Insect metamorphosis is regulated by hormones.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings



- Two systems coordinate communication throughout the body: the endocrine system and the nervous system.
- The endocrine system secretes hormones that coordinate slower but longer-acting responses including reproduction, development, energy metabolism, growth, and behavior.
- The nervous system conveys high-speed electrical signals along specialized cells called neurons; these signals regulate other cells.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Hormones and other signaling molecules bind to target receptors, triggering specific response pathways


- Chemical signals bind to receptor proteins on target cells.
- Only target cells respond to the signal.

Types of Secreted Signaling Molecules

- Secreted chemical signals include
 - Hormones
 - Local regulators
 - Neurotransmitters
 - Neurohormones
 - Pheromones

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Intercellular communication by secreted molecules




Hormones


- Endocrine signals (hormones) are secreted into extracellular fluids and travel via the bloodstream.
- Endocrine glands are ductless and secrete hormones directly into surrounding fluid.
- Hormones mediate responses to environmental stimuli and regulate growth, development, and reproduction.
- Exocrine glands have ducts and secrete substances onto body surfaces or into body cavities (for example, tear ducts).

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Local Regulators = Short Distance Chemical Signals

- **Local regulators** are chemical signals that travel over *short distances* by diffusion.
- Local regulators help regulate blood pressure, nervous system function, and reproduction.
- Local regulators are divided into two types:
 - Paracrine signals act on cells near the secreting cell.
 - Autocrine signals act on the secreting cell itself.





Neurotransmitters and Neurohormones

- Neurons (nerve cells) contact target cells at synapses.
- At synapses, neurons often secrete chemical signals called neurotransmitters that diffuse a short distance to bind to receptors on the target cell. Neurotransmitters play a role in sensation, memory, cognition, and movement.
- **Neurohormones** are a class of hormones that originate *from neurons in the brain* and diffuse through the *bloodstream*.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Pheromones

- Pheromones are chemical signals that are released from the body and used to communicate with other individuals in the species.
- Pheromones mark trails to food sources, warn of predators, and attract potential mates.

Chemical Classes of Hormones


- Three major classes of molecules function as hormones in vertebrates:
 - Polypeptides (proteins and peptides)
 - Amines derived from amino acids
 - Steroid hormones

Polypeptides and amines are water-soluble.

Steroids are lipid-soluble.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Hormones differ in form and solubility



- Lipid-soluble hormones (steroid hormones) pass easily through cell membranes.
- Water-soluble hormones (polypeptides and amines) do not pass through the cell membrane.
- The solubility of a hormone correlates with the location of receptors inside or on the surface of target cells.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Cellular Response Pathways

- Water and lipid soluble hormones differ in their paths through a body.
- Water-soluble hormones are secreted by exocytosis, travel freely in the bloodstream, and bind to cell-surface receptors.
- Lipid-soluble hormones diffuse across cell membranes, travel in the bloodstream bound to transport proteins, and diffuse through the plasma membrane of target cells.

signal transduction pathway

- **Signaling** by any of these hormones involves three key events:
 - Reception
 - Signal transduction
 - Response
- Binding of a hormone to its receptor initiates a signal transduction pathway leading to responses in the cytoplasm, enzyme activation, or a change in gene expression.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings


Receptor location varies with hormone type

Signal receptor

TARGET CELL


(a) NUCLEUS (b)


Receptor location varies with hormone type



Pathway for Water-Soluble Hormones

- The hormone epinephrine has multiple effects in mediating the body's response to short-term stress.
- Epinephrine binds to receptors on the plasma membrane of liver cells.
- This triggers the release of messenger molecules that activate enzymes and result in the release of glucose into the bloodstream.






Pathway for Lipid-Soluble Hormones


- The response to a lipid-soluble hormone is usually a change in gene expression.
- Steroids, thyroid hormones, and the hormonal form of vitamin D enter target cells and bind to protein receptors in the cytoplasm or nucleus.
- Protein-receptor complexes then act as transcription factors in the nucleus, regulating transcription of specific genes.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Multiple Effects of Hormones


- The same hormone may have different effects on target cells that have
 - Different receptors for the hormone
 - Different signal transduction pathways
 - Different proteins for carrying out the response.
- A hormone can also have different effects in different species.





Signaling by Local Regulators

- In paracrine signaling, nonhormonal chemical signals called local regulators elicit responses in nearby target cells.
- Types of local regulators:
 - Cytokines and growth factors
 - Nitric oxide (NO)
 - Prostaglandins help regulate aggregation of platelets, an early step in formation of blood clots.




Simple Hormone Pathways


- Negative feedback and antagonistic hormone pairs are common features of the endocrine system.
- Hormones are assembled into regulatory pathways.
- Hormones are released from an endocrine cell, travel through the bloodstream, and interact with the receptor or a target cell to cause a physiological response.


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings


Insulin and Glucagon: Control of Blood Glucose

- A negative feedback loop inhibits a response by reducing the initial stimulus.
- Negative feedback reverses a trend to regulate many hormonal pathways involved in homeostasis.
- Insulin and glucagon are antagonistic hormones that help maintain glucose homeostasis.
- The pancreas has endocrine cells called islets of Langerhans with alpha cells that produce glucagon and beta cells that produce insulin.









Target Tissues for Insulin and Glucagon

- Insulin reduces blood glucose levels by
 - Promoting the cellular uptake of glucose
 - Slowing glycogen breakdown in the liver
 - Promoting fat storage.
- Glucagon increases blood glucose levels by
 - Stimulating conversion of glycogen to glucose in the liver
 - Stimulating breakdown of fat and protein into glucose.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings


Diabetes Mellitus

- Diabetes mellitus is an endocrine disorder caused by a deficiency of insulin or a decreased response to insulin in target tissues.
- It is marked by elevated blood glucose levels.
- Type I diabetes mellitus (insulin-dependent) is an autoimmune disorder in which the immune system destroys pancreatic beta cells.
- Type II diabetes mellitus (non-insulin-dependent) involves insulin deficiency or reduced response of target cells due to change in insulin receptors.

The endocrine and nervous systems act individually and together in regulating animal physiology

• Signals from the nervous system initiate and regulate endocrine signals.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings



Coordination of Endocrine and Nervous Systems in Invertebrates


- In insects, *molting and development are* controlled by a combination of hormones:
 - A brain hormone stimulates release of ecdysone from the prothoracic glands
 - Juvenile hormone promotes retention of larval characteristics
 - Ecdysone promotes molting (in the presence of juvenile hormone) and development (in the absence of juvenile hormone) of adult characteristics


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings


Coordination of Endocrine and Nervous Systems in Vertebrates

- The hypothalamus receives information from the nervous system and initiates responses through the endocrine system.
- Attached to the hypothalamus is the pituitary gland composed of the posterior pituitary and anterior pituitary.

- The **posterior pituitary** stores and secretes hormones that are made in the hypothalamus
- The anterior pituitary makes and releases hormones under regulation of the hypothalamus







Posterior Pituitary Hormones


- Oxytocin induces uterine contractions and the release of milk
- Suckling sends a message to the hypothalamus via the nervous system to release oxytocin, which further stimulates the milk glands
- This is an example of positive feedback, where the stimulus leads to an even greater response
- Antidiuretic hormone (ADH) enhances water reabsorption in the kidneys

Converget © 2009 Beam on Education Inc., publishing as Beam on Benjamin Cummings

A simple neurohormone pathway Sensory neuron Hypothalamus/ posterior pituitary Response Copyret 2020 Peace (Sacalos vs. publicage on horse Response Copyret 2020 Peace (Sacalos vs. pu


Anterior Pituitary Hormones


- Hormone production in the anterior pituitary is controlled by releasing and inhibiting hormones from the hypothalamus
- For example, the production of thyrotropin releasing hormone (TRH) in the hypothalamus stimulates secretion of the thyroid stimulating hormone (TSH) from the anterior pituitary




Hormone Cascade Pathways

- A hormone can stimulate the release of a series of other hormones, the last of which activates a nonendocrine target cell; this is called a hormone cascade pathway.
- The release of thyroid hormone results from a hormone cascade pathway involving the hypothalamus, anterior pituitary, and thyroid gland.
- Hormone cascade pathways are usually regulated by negative feedback.







Tropic Hormones

- A **tropic hormone** regulates the function of endocrine cells or glands.
- The four strictly tropic hormones are:
 - Thyroid-stimulating hormone (TSH)
 - Follicle-stimulating hormone (FSH)
 - Luteinizing hormone (LH)
 - Adrenocorticotropic hormone (ACTH)

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Growth Hormone

- Growth hormone (GH) is secreted by the anterior pituitary gland and has tropic and nontropic actions.
- It promotes growth directly and has diverse metabolic effects.
- It stimulates production of growth factors.
- An excess of GH can cause gigantism, while a lack of GH can cause dwarfism.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Nontropic Hormones - target nonendocrine tissues.

- Nontropic hormones produced by the anterior pituitary are:
 - Prolactin (PRL)
 - Melanocyte-stimulating hormone (MSH)
- Prolactin stimulates lactation in mammals but has diverse effects in different vertebrates.
- MSH influences skin pigmentation in some vertebrates and fat metabolism in mammals.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Endocrine glands respond to diverse stimuli in regulating metabolism, homeostasis, development, and behavior

• Endocrine signaling regulates metabolism, homeostasis, development, and behavior.

Thyroid Hormone: Control of Metabolism and Development

- The **thyroid gland** consists of two lobes on the ventral surface of the trachea.
- It produces two iodine-containing hormones:
 triiodothyronine (T₃) and thyroxine (T₄).
- Proper thyroid function requires dietary iodine for thyroid hormone production.


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Parathyroid Hormone and Vitamin D: Control of Blood Calcium

- Two antagonistic hormones regulate the homeostasis of calcium (Ca²⁺) in the blood of mammals
 - Parathyroid hormone (PTH) is released by the parathyroid glands
 - Calcitonin is released by the thyroid gland

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

- Thyroid hormones stimulate metabolism and influence development and maturation.
- Hyperthyroidism, excessive secretion of thyroid hormones, causes high body temperature, weight loss, irritability, and high blood pressure.
- Graves' disease is a form of hyperthyroidism in humans.
- Hypothyroidism, low secretion of thyroid hormones, causes weight gain, lethargy, and intolerance to cold.



• PTH increases the level of blood Ca2+

- It releases Ca²⁺ from bone and stimulates reabsorption of Ca²⁺ in the kidneys
- It also has an indirect effect, stimulating the kidneys to activate vitamin D, which promotes intestinal uptake of Ca²⁺ from food
- Calcitonin decreases the level of blood Ca²⁺
 - It stimulates Ca²⁺ deposition in bones and secretion by kidneys

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Catecholamines from the Adrenal Medulla

- The adrenal medulla secretes epinephrine (adrenaline) and norepinephrine (noradrenaline).
- These hormones are members of a class of compounds called catecholamines.
- They are secreted in response to stressactivated impulses from the nervous system.
- They mediate various fight-or-flight responses.


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings


Adrenal Hormones: Response to Stress


- The adrenal glands are adjacent to the kidneys.
- Each adrenal gland actually consists of two glands: the adrenal medulla (inner portion) and adrenal cortex (outer portion).

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

- Epinephrine and norepinephrine
 - Trigger the release of glucose and fatty acids into the blood
 - Increase oxygen delivery to body cells
 - Direct blood toward heart, brain, and skeletal muscles, and away from skin, digestive system, and kidneys.
- The release of epinephrine and norepinephrine occurs in response to nerve signals from the hypothalamus.








Steroid Hormones from the Adrenal Cortex

- The adrenal cortex releases a family of steroids called **corticosteroids** in response to stress.
- These hormones are triggered by a hormone cascade pathway via the hypothalamus and anterior pituitary.
- Humans produce two types of corticosteroids: glucocorticoids and mineralocorticoids.



- Glucocorticoids, such as cortisol, influence glucose metabolism and the immune system.
- Mineralocorticoids, such as aldosterone, affect salt and water balance.
- The adrenal cortex also produces small amounts of steroid hormones that function as sex hormones.

Gonadal Sex Hormones

- The gonads = testes and ovaries, produce most of the sex hormones: androgens, estrogens, and progestins.
- All three sex hormones are found in both males and females, but in different amounts.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

- The testes primarily synthesize androgens, mainly testosterone, which stimulate development and maintenance of the male reproductive system and male secondary sex characteristics.
- Testosterone causes an increase in muscle and bone mass and is often taken as a supplement to cause muscle growth, which carries health risks.

- Estrogens, made in the ovary, most importantly estradiol, are responsible for maintenance of the female reproductive system and the development of female secondary sex characteristics.
- In mammals, progestins, which include progesterone, are primarily involved in preparing and maintaining the uterus.
- Synthesis of the sex hormones is controlled by FSH and LH from the anterior pituitary.

Signal Transduction Pathway Low blood glucose Pancreas alpha cells secretes glucagon Pancreas alpha cells secretes glucagon Cayyore 2000 Peace Canage of Pancreas alpha cells secretes glucagon

Pineal Gland - Melatonin and Biorhythms

- The pineal gland, located in the brain, secretes melatonin.
- Light/dark cycles control release of melatonin.
- Primary functions of melatonin appear to relate to biological rhythms associated with reproduction.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

You should now be able to:

- 1. Distinguish between the following pairs of terms: hormones and local regulators, paracrine and autocrine signals.
- 2. Describe the evidence that steroid hormones have intracellular receptors, while watersoluble hormones have cell-surface receptors.
- 3. Explain how the antagonistic hormones insulin and glucagon regulate carbohydrate metabolism.
- 4. Distinguish between type 1 and type 2 diabetes.

- 5. Explain how the hypothalamus and the pituitary glands interact and how they coordinate the endocrine system.
- 6. Explain the role of tropic hormones in coordinating endocrine signaling throughout the body.
- 7. List and describe the functions of hormones released by the following: anterior and posterior pituitary lobes, thyroid glands, parathyroid glands, adrenal medulla, adrenal cortex, gonads, pineal gland.