

# MTH 102 Calculus -II

Dr. Gizem SEYHAN ÖZTEPE

Topic-11-Functions of Several Variables

The volume  $V$  of a circular cylinder depends on its radius  $r$  and its height  $h$ . In fact, we know that  $V = \pi r^2 h$ . We say that  $V$  is a function of  $r$  and  $h$ , and we write  $V(r, h) = \pi r^2 h$ .

## Definition

A function  $f$  of two variables is a rule that assigns to each ordered pair of real numbers  $(x, y)$  in a set  $D$  a unique real number denoted by  $f(x, y)$ . The set  $D$  is the domain of  $f$  and its range is the set of values that  $f$  takes on, that is,  $\{f(x, y) : (x, y) \in D\}$ .

We often write  $z = f(x, y)$ . So,  $x$  and  $y$  are independent variables;  $z$  is a dependent variable.





Figure 1 shows the domain and range of two variable functions.


FIGURE 1

- ▶ The set of points in the plane where a function  $f(x, y)$  has a constant value  $f(x, y) = c$  is called a **level curve** of  $f$ .
- ▶ The set of all points  $(x, y, f(x, y))$  in space is called the **graph** of  $f$ . The graph of  $f$  is also called the surface  $z = f(x, y)$ .




This figure shows the graph of a surface


You can see the graph of several functions as follows:




(a)  $f(x,y) = (x^2 + 3y^2)e^{-x^2-y^2}$



(b)  $f(x,y) = (x^2 + 3y^2)e^{-x^2-y^2}$



(c)  $f(x, y) = \sin x + \sin y$



(d)  $f(x, y) = \frac{\sin x \sin y}{xy}$

# Limits and Continuity

**Definition (Limit)** Let  $f$  be a function of two variables whose domain  $D$  includes points arbitrarily close to  $(a, b)$ . Then we say that the limit of  $f(x, y)$  as  $(x, y)$  approaches  $(a, b)$  is  $L$  and we write

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$$

if for every number  $\varepsilon > 0$  there is a corresponding number  $\delta > 0$  such that if  $(x, y) \in D$  and  $0 < \sqrt{(x - a)^2 + (y - b)^2} < \delta$  then  $|f(x, y) - L| < \varepsilon$


If  $f(x, y) \rightarrow L_1$  as  $(x, y) \rightarrow (a, b)$  along a path  $C_1$  and  $f(x, y) \rightarrow L_2$  as  $(x, y) \rightarrow (a, b)$  along a path  $C_2$ , where  $L_1 \neq L_2$ , then  $\lim_{(x,y) \rightarrow (a,b)} f(x, y)$  does not exist.

### Definition (Continuity):


A function  $f$  of two variables is called continuous at  $(a, b)$  if

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = f(a, b)$$

We say  $f$  is continuous on  $D$  if  $f$  is continuous at every point  $(a, b)$  in  $D$ .



Graph of a continuous function



Graph of a discontinuous function