



European Code against Cancer 4th Edition: Diet and cancer[☆]



Teresa Norat^a, Chiara Scoccianti^b, Marie-Christine Boutron-Ruault^c, Annie Anderson^d, Franco Berrino^e, Michele Cecchini^{f,1}, Carolina Espina^b, Tim Key^g, Michael Leitzmann^h, Hilary Powersⁱ, Martin Wiseman^j, Isabelle Romieu^{b,*}

^a Department of Epidemiology and Biostatistics, School of Public Health Imperial College London, St Mary's Campus, London W2 1PG, United Kingdom

^b International Agency for Research on Cancer (IARC), 150 Cours Albert Thomas, 69372 Lyon Cedex 08, France

^c Institut Gustave Roussy, 114 rue Edouard Vaillant, 94805 Villejuif, France

^d Centre for Research into Cancer Prevention & Screening, Level 7, Mailbox 7, Ninewells Hospital & Medical School, Dundee DD1 9SY, Scotland, United Kingdom

^e Fondazione IRCCS Istituto Nazionale dei Tumori, 1 via Venezian, 20133 Milan, Italy

^f Health Policy Analyst OECD, 2 rue André Pascal, 75775 Paris Cedex 16, France

^g Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, United Kingdom

^h Department of Epidemiology and Preventive Medicine, University Hospital Regensburg, 93042 Regensburg, Germany

ⁱ Human Nutrition Unit, The Medical School, Beech Hill Road, Sheffield S10 2RX, United Kingdom

^j World Cancer Research Fund International, Second Floor, 22 Bedford Square, London WC1B 3HH, United Kingdom

ARTICLE INFO

Article history:

Received 2 July 2014

Received in revised form 19 December 2014

Accepted 30 December 2014

Available online 9 July 2015

Keywords:

Primary prevention

Europe

Neoplasms/aetiology/*prevention & control

*Diet

Sodium chloride

Dietary/administration & dosage/adverse effects

*Fruit

*Vegetables

Meat/adverse effects

ABSTRACT

Lifestyle factors, including diet, have long been recognised as potentially important determinants of cancer risk. In addition to the significant role diet plays in affecting body fatness, a risk factor for several cancers, experimental studies have indicated that diet may influence the cancer process in several ways. Prospective studies have shown that dietary patterns characterised by higher intakes of fruits, vegetables, and whole-grain foods, and lower intakes of red and processed meats and salt, are related to reduced risks of death and cancer, and that a healthy diet can improve overall survival after diagnosis of breast and colorectal cancers. There is evidence that high intakes of fruit and vegetables may reduce the risk of cancers of the aerodigestive tract, and the evidence that dietary fibre protects against colorectal cancer is convincing. Red and processed meats increase the risk of colorectal cancer. Diets rich in high-calorie foods, such as fatty and sugary foods, may lead to increased calorie intake, thereby promoting obesity and leading to an increased risk of cancer. There is some evidence that sugary drinks are related to an increased risk of pancreatic cancer.

Taking this evidence into account, the 4th edition of the European Code against Cancer recommends that people have a healthy diet to reduce their risk of cancer: they should eat plenty of whole grains, pulses, vegetables and fruits; limit high-calorie foods (foods high in sugar or fat); avoid sugary drinks and processed meat; and limit red meat and foods high in salt.

© 2015 International Agency for Research on Cancer; Licensee ELSEVIER Ltd <https://creativecommons.org/licenses/by-nc-nd/3.0/igo/>

[☆] This is an Open Access article published under the CC BY NC ND 3.0 IGO license which permits users to download and share the article for non-commercial purposes, so long as the article is reproduced in the whole without changes, and provided the original source is properly cited. This article shall not be used or reproduced in association with the promotion of commercial products, services or any entity. There should be no suggestion that IARC endorses any specific organisation, products or services. The use of the IARC logo is not permitted. This notice should be preserved along with the article's original URL.

Abbreviations: BMI, body mass index; ECAC, European Code against Cancer; EPIC, European prospective study; WCRF/AICR, World Cancer Research Fund/American Institute for Cancer Research; WHEL, Women's Healthy Eating and Living Trial; WINS, Women's Intervention Nutrition Study.

* Corresponding author at: IARC European Code against Cancer Secretariat, 150 Cours Albert Thomas, F-69372 Lyon Cedex 08, France. Tel.: +33 04 72 73 84 85.

E-mail address: secretariat-cancer-code-europe@iarc.fr (I. Romieu).

¹ The views expressed are those of the author and not necessarily those of the OECD, or its member countries.

1. Introduction

Lifestyle factors, including diet, have long been recognised as potentially important determinants of cancer risk. In 1981, Doll and Peto estimated that 35% of cancer deaths might be avoidable by changes in diet and tackling obesity [1]. Ecological studies showing significant correlations between dietary habits and cancer incidence and mortality [2], and the large global variations in cancer incidence and mortality rates, along with rapid changes in cancer rates among migrant populations [3], stimulated further research on the role of diet in cancer prevention.

The most extensive review of the existing evidence on diet and cancer is the 2007 World Cancer Research Fund/American Institute for Cancer Research (WCRF/AICR) [4] report and its subsequent

updates [5]. Lifestyle recommendations for cancer prevention were drawn up on the basis of nutrition-related factors judged to be convincingly or probably causally related to cancer, according to predefined criteria for judging the strength of the evidence regarding causality [6] (Table 1). According to the recommendations, a healthy diet for cancer prevention is a diet (1) that allows a person to be as lean as possible without being underweight; (2) is rich in fruits, vegetables, whole grains and pulses; (3) contains low amounts of red meat; (4) does not contain processed meats; and (5) limits salt intake. In addition, a healthy diet is characterised by the avoidance of sugary drinks and limited intake of calorie-rich foods, thereby contributing to achieving and maintaining a healthy weight. A healthy diet also limits consumption of alcoholic drinks. The preventability of several cancers in the UK through healthy eating, being physically active, and maintaining a healthy weight as recommended by WCRF/AICR was estimated to range from 5% to 34% [7] (Fig. 1).

1.1. Main mechanisms involving nutritional factors in cancer development

Excessive caloric intake results in weight gain and ultimately obesity, which in turn is associated with an increased risk of several cancers. Many experimental studies have shown that calorie restriction suppresses the carcinogenic process [8]. The biological mechanisms linking adiposity and cancer risk include hyperinsulinaemia and insulin resistance, up-regulation of insulin-like growth factors, modification of the metabolism of sex hormones, chronic inflammation, changes in production of adipokines and vascular growth factors by adipose tissue, oxidative stress, and alterations in immune function [9].

In addition to its caloric content, diet may influence cellular processes and lead to the accumulation of the eight hallmarks of cancer cells: self-sufficiency in growth signals, insensitivity to anti-growth signals, limitless replicative potential, evasion of apoptosis, sustained angiogenesis, reprogramming of energy metabolism, evasion of immune destruction, tissue evasion and metastasis [10].

The links between diet and cancer are complex. Thousands of dietary components are consumed each day; a typical diet may provide more than 25,000 bioactive food constituents, and the amounts of bioactive components within a particular food may widely vary [11]. Each bioactive food constituent has the potential to modify multiple aspects of the cancer process, alone or in combination with several micronutrients, and the quantity, timing, and duration of exposure modulate the cell response. Thus, it is not possible to ascribe a causal effect to specific compounds; it is more likely that the effect results from a combination of influences on several pathways involved in carcinogenesis.

A growing body of evidence indicates that lowering the energy density (the amount of energy in a particular weight of food) of diets can reduce caloric intake (reviewed by Rolls [12]). Energy-dense diets contain less fibre-rich foods, and are usually high in fats, processed starch, and added sugars.

Trans fatty acids are used in industrially processed sweet and salty foods, such as chocolate bars, candies, biscuits, cakes, crackers, industrial bread, and packaged snacks. The array of potentially harmful effects of industrial trans fatty acids is wide, and include alterations in metabolic and signalling pathways, higher circulating levels of lipid, systemic inflammation, endothelial dysfunction, and possibly increased visceral adiposity, body weight and insulin resistance [13]. There is evidence that trans fatty acids could be associated with an increased risk of breast cancer [14,15], non-aggressive prostate cancer [16], gastric adenocarcinomas [17] and colorectal adenomas [18]. The World Health Organisation recommends eliminating the use of trans fatty acids in food processing for cardiovascular disease prevention [19].

Table 1

2007 World Cancer Research Fund – American Institute for Cancer Research Personal Recommendations for Cancer Prevention.

Recommendations

Body fatness

Be as lean as possible within the normal range of body weight

Personal recommendations

Ensure that body weight throughout childhood and adolescent growth projects towards the lower end of the normal BMI range at age 21

Maintain body weight within the normal range from age 21

Avoid weight gain and increases in waist circumference throughout adulthood

Physical activity

Be physically active as part of everyday life

Personal recommendations

Be moderately physically active, equivalent to brisk walking, for at least 30 min every day

As fitness improves, aim for ≥ 60 min of moderate or ≥ 30 min of vigorous physical activity every day

Limit sedentary habits such as watching television

Foods and drinks that promote weight gain

Limit consumption of energy-dense foods and avoid sugary drinks

Personal recommendations

Consume energy-dense foods sparingly

Avoid sugary drinks

Consume fast foods sparingly, if at all

Plant foods

Eat foods mostly of plant origin

Personal recommendations

Eat at least five portions/servings (at least 400 g or 14 oz) of a variety of non-starchy vegetables and fruit every day

Eat relatively unprocessed cereals (grains) and/or pulses (legumes) with every meal

Limit refined starchy foods

People who consume starchy roots or tubers as staples should also to ensure sufficient intake of non-starchy vegetables, fruit, and pulses

Animal foods

Limit intake of red meat and avoid processed meat

Personal recommendations

People who eat red meat to consume <500 g (18 oz) a week, very little – if any – to be processed

Alcoholic drinks

Limit alcoholic drinks

Personal recommendations

If alcoholic drinks are consumed, limit consumption to no more than two drinks a day for men and one drink a day for women

Preservation, processing, preparation

Limit consumption of salt

Avoid mouldy cereals (grains) or pulses (legumes)

Personal recommendations

Avoid salt-preserved, salted, or salty foods; preserve foods without using salt.

Limit consumption of processed foods with added salt to ensure an intake of <6 g (2.4 g sodium) a day

Do not eat mouldy cereals (grains) or pulses (legumes)

Dietary supplements

Aim to meet nutritional needs through diet alone

Personal recommendations

Dietary supplements are not recommended for cancer prevention

Breastfeeding

Mothers to breastfeed; children to be breastfed

Personal recommendations

Aim to breastfeed infants exclusively up to 6 months and continue with complementary feeding thereafter

Cancer survivors

Follow the recommendations for cancer prevention

Personal recommendations

All cancer survivors should receive nutritional care from an appropriately trained professional.

If able to do so, and unless otherwise advised, aim to follow the recommendations for diet, healthy weight, and physical activity

Source: Food, Nutrition, Physical Activity and the Prevention of Cancer: a Global Perspective, 2007 [4].

Notes:

Normal (or healthy) weight correspond to a body mass index between 18.5 and 24.9 kg/m². Examples of light physical activity are walking slowly, light gardening, housework; examples of moderate activities are walking briskly, cycling, dancing, swimming and vigorous activities (running, tennis and football).

Energy-dense foods are defined as those with an energy content of more than about 225–275 kcal/g.

Sugary drinks refer to drinks with added sugars. Fruit juices should also be limited.

“Fast foods” refer to readily available convenience foods that tend to be energy-dense and are frequently consumed in large portions.




Fig. 1. Cancer preventability estimates for the United Kingdom for dietary factors judged to be convincingly or probably related to the risk of some cancers. The figure shows the percentage of cancer cases preventable by adequate diet in the United Kingdom through adequate intakes of dietary factors considered probably or convincingly related to the risk of some cancers [4,5]. Processed meats and red meats convincingly increase the risk of colorectal cancer; foods rich in fibre convincingly decrease the risk of colorectal cancers; salt, salted and salty foods probably increase the risk of stomach cancer; and fruits and vegetables probably decrease the risk of cancers of the mouth, pharynx and larynx, oesophagus and stomach.

Source: Figure based on data from the Diet and Cancer Report website [7].

Diets high in sugars may promote carcinogenesis by increasing insulin production, increasing oxidative stress [20], or promoting weight gain [21]. In metabolic syndrome, foods rich in sugars are likely to interfere with levels of blood glucose and/or triglycerides, either directly or through insulin and other hormones [22], while sugar-sweetened soft drinks play a role in the development of obesity (reviewed by Hu et al. [21]).

Some food compounds are direct mutagens and carcinogens. Many nitroso compounds are mutagenic and potentially carcinogenic. Haem iron, which is abundant in red meat but not in white meat, promotes colorectal carcinogenesis through its catalytic activity in the formation of nitroso compounds and of lipid oxidation end products such as 4-hydroxynonenal [23]. In processed red meat, haem iron is nitrosylated, because curing salt contains nitrate or nitrite; there is evidence that nitrosylated haem iron promotes carcinogenesis at 5–6 times lower doses than non-nitrosylated haem iron. Experimental studies show that calcium salts, chlorophyll, vitamin C, and several polyphenols may reduce the deleterious effects of haem (reviewed by Bastide et al. [24]). Heterocyclic amines and polycyclic aromatic hydrocarbons are other potential carcinogens that are formed in meats when cooked at high temperatures for a long time, or exposed to a direct flame [25,26]; this is not limited to red and processed meats. Various single-nucleotide polymorphisms involved in the metabolism of these potential carcinogens may modulate the association of carcinogens formed in meat with cancer risk [27–29].

Dietary fibre from plant foods stimulates bacterial anaerobic fermentation in the large bowel, leading to the production of short-chain fatty acids: acetate, propionate, and butyrate. In cell lines, butyrate reduces cell proliferation and induces apoptosis. Dietary fibre may protect against colorectal cancer by reducing the contact between the intestinal content and the mucosa, and may interfere with the enterohepatic circulation of oestrogens [30]. Some studies have shown reductions in circulating oestrogen and androstenedione levels with high fibre intake [31,32], and in experimental studies in mice soluble fibre reduced mammary tumour growth, angiogenesis and metastasis [33]. In addition, fibre-rich foods are important sources of phyto-oestrogens, oestrogen-like plant compounds that may interact with and modulate the activity of

oestrogen receptors [34,35], thereby modulating the risk of hormone-dependent cancers (especially breast cancer) [36]. Polyphenols from plant foods have multiple biological effects, including scavenging of oxidative agents, anti-inflammatory and detoxifying actions, inhibition of platelet aggregation, and antimicrobial activity [37,38].

Vitamins and minerals such as carotenoids, folate, vitamins C, D, E and B6, selenium, and phytochemicals might reduce cancer risk through preventing oxidative damage, inhibiting cell proliferation, inducing cell-cycle arrest, maintaining DNA methylation, and/or modulating steroid hormone concentrations and hormonal metabolism (e.g., via phyto-oestrogen contained in pulses) [39].

Diallyl disulphide from garlic and other allium vegetables, and sulphoraphane, a glucosinolate from cruciferous vegetables, can behave as histone deacetylase inhibitors and act to maintain DNA stability or enhance transcription [40]. Naturally occurring bioactive compounds such as curcumin, resveratrol, and isothiocyanates have potential antioxidant and/or anti-inflammatory and anti-carcinogenic activities (reviewed by Chung et al. [41]).

Sodium chloride is a food preservative used for increasing the safety and shelf life of processed foods. In animal experiments, salt intake facilitates gastric colonisation by *Helicobacter pylori*, one of the main predisposing factors for stomach cancer development, and induces mucosal damage [42], potentially leading to chronic atrophic gastritis. Salt intake may also promote or enhance the effect of nitroso compounds and other carcinogens (reviewed by D'Elia et al. [43]).

1.2. Overview of the main European dietary patterns

As a consequence of the convergence of factors that influence food choices, differences in dietary patterns across European countries are becoming less marked. However, data from food availability at the national and household levels, and dietary surveys on individual food consumption (European Nutrition and Health Report 2009 [44]), show that regional differences still exist (Fig. 2). Similar findings had previously been reported in large population studies that collected data on food and beverage intakes using detailed dietary questionnaires [45]. On average, the




Fig. 2. Mean household availability of selected food and beverages in European regions. The figure shows the mean quantity (g/day) of selected foods available for each person in a household in European regions, assuming an equal distribution of foods across household members. Soft drinks are in ml/person/day. Foods and beverages consumed in restaurants, canteens or similar establishments were not consistently available. The countries included in the survey were Finland, Latvia, Norway, Sweden (North); Austria, Germany, Hungary, Poland, Slovenia (Central and Eastern), Belgium, France, Ireland, Luxembourg, United Kingdom (West); Cyprus, Greece, Italy, Spain, Portugal (South). The year of data collection ranged from 1991 to 2005, depending on the country.

Source: Figure based on data from Elmadfa et al. [44].

most marked difference is the North–South gradient of fruits, vegetables, cereals and pulses availability. Southern Europe is characterised by a lower availability of processed meats. The observed pattern is similar to the “Mediterranean” pattern, which is characterised by high intakes of vegetables, grains, dried beans, olive oil, seafood and fruits, and by moderate wine intake during meals.

The highest availability of milk and dairy-based products is found in Northern Europe, which, as in Central and Eastern Europe, is characterised by a higher household availability of processed meats. Sugary drinks are more available in Western European households [46]. Overall, higher educational level in the household is correlated with higher availability of fish, fruits, and vegetables. Lower socioeconomic status is associated with more frequent and greater availability of soft drinks [47].

2. Evidence of the association of diet and cancer in population studies

The randomised controlled trial is considered the best study design for assessing the effect of an intervention because the processes used to conduct this study type minimise the possibility that confounding factors could influence the results. However, randomised controlled trials are only justified when there is considerable evidence from animal, *in vitro* and observational studies that the intervention could have a beneficial effect. Testing potentially harmful interventions would not be ethical. Randomised controlled trials are also limited by a number of methodological issues, including the difficulty of incorporating whole-diet interventions into a randomised controlled trial design and in a not-blinded way, the compliance of study participants with dietary changes over a long period of time, and determination of the adequate nutrient dose or dietary intervention for a beneficial effect to occur.

Randomised controlled trials of nutrition and cancer are rare compared with observational studies. For less frequent cancers, the evidence from observational studies comes mainly from case–control studies. In these studies, dietary intake is assessed using questionnaires after cancer diagnosis, and they are more prone to recall and selection biases than are prospective observational studies. For more frequent cancers, most of the existing data on the relationship between diet and cancer risk comes from prospective observational studies, in which dietary assessment takes place before cancer diagnosis. The identification of new cancer cases during follow-up includes record linkage with cancer and mortality registries, self-reporting by study participants with further confirmation through medical records, or a combination of methods to minimise losses to follow-up and increase data reliability. Prospective studies are expensive, and several years are needed before the number of cancer cases required for analyses can be documented.

2.1. Fruits, vegetables, pulses, and whole-grain foods

A potentially protective effect of fruits and vegetables against cancer was supported by evidence from earlier case–control studies [48]. Subsequent data from prospective studies indicate that the association may be restricted to specific cancers and may be weaker than previously observed for some cancers [4,49].

The accumulated evidence supports the idea that low intake of fruits and vegetables is related to the development of cancers of the respiratory and upper digestive tracts (Table 2).

A recently published large consortium of case–control studies confirmed the inverse association between fruits and vegetables and cancers of the mouth, pharynx and larynx [50]. Data from cohort studies, which are less prone to bias than case–control studies, are needed to confirm these findings.

Recent cohort studies are also supportive of an inverse association between fruit intake and oesophageal squamous-cell

Table 2
Estimated associations from meta-analyses, pooled analyses and recent prospective studies for dietary factors judged in the 2007 WCRF/AICR as convincingly or probably causes of or protective against cancer.

Cancer type	Number of studies	Comparison unit	Relative risk	Reference
<i>Dietary fibre</i>				
Colon cancer	16 cohorts	For 10 g/d increase	0.90 (0.86–0.94)	[72]
<i>Vegetables</i>				
Mouth, pharynx, and larynx cancer	22 case-control studies	Highest versus lowest	0.68 (0.51–0.90)	[50]
Oesophageal cancer (squamous cell carcinoma)	5 cohorts	Highest versus lowest	0.80 (0.61–1.07)	[51]
Oesophageal cancer (all)	5 case-control studies	For 50 g/d increase	0.87 (0.72–1.05)	[4]
Stomach cancer	19 cohort studies	Highest versus lowest	0.96 (0.88–1.06)	[56]
Stomach cancer	1 cohort, 7 case-control (Japanese or Korean studies)	Highest versus lowest	0.62 (0.46–0.85)	[134]
	7 cohorts	For 100 g/d increase	0.98 (0.91–1.06)	[4]
<i>Allium vegetables:</i>				
Stomach cancer	2 cohorts, 12 case-control studies	Highest versus lowest allium vegetables intake	0.54 (0.43–0.65)	[63]
<i>Garlic</i>				
Colorectal cancer	2 cohorts	Highest versus lowest	Women: 1.21 (0.94–1.57) Men: 1.00 (0.71–1.42)	[135]
	1 cohort	For each additional clove or “4 shakes” of garlic per week	1.04 (0.99–1.08)	[136]
	1 cohort	Any pills per day previous 10 years versus no use	1.35 (1.01–1.81)	[137]
	2 cohorts	Highest versus lowest	0.72 (0.54–0.96)	[4]
<i>Fruits</i>				
Head and neck cancer	22 case-control	Highest versus lowest	0.52 (0.43–0.62)	[50]
Upper aero-digestive tract cancer	3 cohort studies	Highest versus lowest	0.78 (0.64–0.95)	[138]
Oesophageal cancer (squamous cell carcinoma)	5 cohorts	Highest versus lowest	0.68 (0.55–0.86)	[51]
Oesophageal cancer (all)	8 case-control	For 100 g/d increase	0.56 (0.42–0.74)	[4]
Lung cancer	7 cohorts (European)	Per 1 g/day	0.99 (0.99–0.99)	[139]
	14 cohorts	For one serving/day increase	0.94 (0.90–0.97)	[4]
Stomach cancer	22 cohort studies	Highest versus lowest	0.90 (0.83–0.98)	[56]
Stomach cancer		For 100 g/d increment	0.95 (0.89–1.02)	[4]
<i>Red meat</i>				
Colorectal cancer	10 cohorts	For 100 g/d increment	1.17 (1.02–1.33)	[95]
<i>Processed meat</i>				
Colorectal cancer	9 cohorts	For 50 g/day increment	1.18 (1.10–1.28)	[95]
<i>Salt intake</i>				
Stomach cancer	4 cohorts and 7 case-control studies	Highest versus lowest salt intake	1.22 (1.17–1.27)	[100]
	7 studies	Salt intake from salty food	2.41 (2.08–2.78)	[101]
	7 cohorts	Highest versus lowest salt intake	1.68 (1.17–2.41)	[4]
	2 cohorts	For 1 g/d increment salt intake	1.08 (1.00–1.17)	
<i>Foods rich in salt</i>				
Stomach cancer	10 cohorts	Use versus never or occasional consumption of pickle vegetables	1.32 (1.10–1.59)	[140]
<i>Pickled food</i>				
Stomach cancer	11 cohorts	Highest versus lowest	1.27 (1.09–1.49)	[101]
<i>Salted fish</i>				
Stomach cancer	13 cohorts	Highest versus lowest	1.24 (1.03–1.50)	[101]
<i>Salted and salty foods</i>				
Stomach cancer	4 cohorts	For 1 serving/day	1.32 (0.90–1.95)	[4]
<i>Soft sugary beverages</i>				
Pancreatic cancer	11 cohorts	Highest versus lowest	1.12 (0.99–1.27)	[89]
<i>Fructose</i>				
Pancreatic cancer	6 cohorts	For 25 g/day increment	1.22 (1.08–1.37)	[90]

carcinoma (summarised by Liu et al., 2013 [51]); the evidence of an association with oesophageal adenocarcinoma is weaker. With respect to specific types of fruits, no significant associations between intake of citrus fruits and oesophageal cancer were reported in two Asian cohorts [52,53], and the association between oesophageal cancer and dietary vitamin C was not confirmed by two recent cohort studies [54,55]. Oesophageal cancer was shown

to be inversely associated with intake of vegetables in a meta-analysis of cohort studies [51], although no statistical significance was observed.

The evidence for fruit intake and stomach cancer has been confirmed in a meta-analysis of 22 cohort studies showing that individuals with lower fruit intake are at higher risk of stomach cancer [56]. The association between stomach cancer and citrus

fruit intake was also confirmed in a meta-analysis of 12 case-control studies and two cohort studies [57]. These results were not confirmed in three recent reports from cohort studies [54,58,59]. The possibility that errors in diet measurement have obscured the association between citrus fruit intake and gastric cancer was suggested by a large multinational European prospective study (EPIC). This study reported an inverse association between plasma levels of vitamin C and gastric cancer risk, whereas no association was observed with dietary vitamin C [60]. Moreover, a meta-analysis of three cohort studies showed a 30% lower risk of stomach cancer for the highest compared to the lowest levels of pre-diagnosis plasma vitamin C [61].

For vegetables, no association with stomach cancer was observed in a large European prospective study (EPIC) [58], other cohort studies [54,59,62], and in a meta-analysis of 22 cohort studies [56]. A protective association of allium vegetables was reported in a meta-analysis of 19 case-control studies and two cohort studies [63]. No association was observed in the large EPIC study [58]. No effect on stomach cancer incidence or mortality was also observed in the Shandong Intervention Trial, China: a double-blind 7-year oral supplementation trial with garlic extract and steam-distilled garlic oil in *Helicobacter pylori* seronegative participants [64].

The inverse association between fruit intake and lung cancer observed in numerous case-control and cohort studies was confirmed in the EPIC study [65]. The NIH-AARP North American cohort study [66] did not find a significant association. As for vegetables, evidence of an association with lung cancer was provided by a meta-analysis of prospective cohort studies [67] and a study in Chinese men showing an inverse association with carotenoid-rich vegetables [68]. In a multi-ethnic cohort, circulating carotenoid levels were inversely related to lung cancer risk in men but not in women [69], raising the issue of potential confounding by smoking. However, the interpretation of these findings is complicated because the metabolism of carotenoids is affected by many hormonal, genetic and lifestyle factors.

The evidence of an association between fruits and vegetables and colorectal cancer was judged as “limited-suggestive” in the 2007 WCRF/AICR report. Evidence for an inverse association was reported in the more recent WCRF/AICR update [70], in which a non-linear relationship was observed, indicating that the benefit from an increase in fruit and vegetable consumption would be limited to people with the lowest intake levels, and that no substantial benefit would occur in people who already have a high intake of high fruits and vegetables. Not included in this update is the large NIH-AARP American cohort study which more recently reported an inverse association between vegetable intake and colon cancer, but no association between fruits and colon cancer [71].

The evidence for a protective role of dietary fibre on colorectal cancer in recent publications is stronger [72], and there is some evidence that it could be stronger for fibre from cereals. Whole-grain foods are rich in fibre, and have also been found to be inversely related to colorectal cancer risk [72]. Pulses are high in fibre and provide substantial amounts of protein, vitamins, and minerals with a relatively low amount of calories [73]. Case-control studies support a beneficial effect of pulses [4], but the data from published cohort studies on pulse intake are limited.

For other cancers, data on the potential beneficial effect of fibre has been accumulating, as shown by meta-analyses of cohort studies on breast cancer [74] and of case-control studies on oesophageal adenocarcinoma [75] and stomach cancer [76].

2.2. High-calorie foods and beverages

High-calorie foods are rich in fats and/or sugars, and often have low nutritional value except from providing energy for the functioning of the body.

In ecological studies, fat intake has been found to be related to higher risk of several cancers. Prospective observational studies have not confirmed the association between dietary fat and colorectal cancer [77], and studies on breast cancer have been inconsistent. There is some evidence that the relationship between dietary fat and breast cancer may have been obscured by imprecise measurement of fat intake [78,79], although this was not confirmed in another large study [80]. Inconsistencies may also have emerged if the intake of specific fats, and not of total fat, is related only to some breast cancer types. In a recent meta-analysis of cohort studies, intake of polyunsaturated fatty acids was weakly associated with higher risk of postmenopausal breast cancer [81], and in the EPIC study, saturated fat intake was related to a higher risk of positive-receptor breast cancer only [82]. In a nested case-control study within the French EPIC cohort that analysed membrane phospholipids, trans fatty acids from industrial sources were associated with increased breast cancer risk [14].

The Women's Health Initiative Dietary Modification Randomised Controlled Trial investigated whether a low-fat dietary pattern intervention could reduce the incidence of breast and other cancers. After approximately 8 years of follow-up, no reduction in the incidence of breast, colorectal, or endometrial cancers was observed; there was only a marginal reduction in the incidence of ovarian cancer [83]. However, women in the study only modestly lowered their fat intake, from 38% to 29%, and no change was seen in blood lipid levels. There was a small weight loss in the early years of the trial in women in the low fat group.

In a large prospective study sugars from beverages – specifically, added fructose from fructose-glucose corn syrup, but not sugars from solid foods – were positively associated with overall mortality but not with the risk of major cancers [84]. Sugar-sweetened beverages raise insulin and glucose levels, and its consumption is associated with obesity and diabetes [85], which have been found to be related to pancreatic cancer risk [86,87]. Intake of sugar-sweetened beverages was not related to pancreatic cancer in a meta-analysis of case-control and cohort studies [88], but a positive association between sugar-sweetened beverages and pancreatic cancer was observed in a pooled analysis of cohort studies [89]. In another recent meta-analysis of cohort studies, a higher risk of pancreatic cancer was observed for higher fructose intake [90]. Fructose has been a component of sugary drinks in North America, but – other than as part of sucrose – not in Europe [91]. Sugary drinks have not been found to be directly related to the risk of colon cancer [92] or other cancers [4], independently of body fatness. Greater consumption of foods with a high glycaemic index was significantly related to increased risks of breast and colorectal cancer in a meta-analysis of 11 and nine prospective studies, respectively [93,94].

2.3. Red and processed meats and cancer

Numerous studies have shown an association between high intake of processed meat (such as ham, bacon, sausages, and hot dogs), red meat (mainly beef, pork or lamb) and colorectal cancer [4,95] (Table 2). Overall, the increased risk associated with processed meat intake was higher than that with unprocessed red meat. Processed meat cured with nitrite contains high concentrations of preformed nitroso compounds and nitrosylated haem iron, and these are potential carcinogens. There is experimental evidence that meats cured with nitrite may increase oxidative DNA damage [93]. Studies on carcinogens formed during the cooking of meat have not been conclusive, perhaps because of interactions with genetic polymorphisms, such as the acetylator phenotypes, and because of the difficulties in assessing dietary carcinogen intake.

Red and processed meat intake has also been found to be related to the risk of stomach [96] and pancreatic cancers [97], and with

higher overall cancer mortality [98,99]. In addition to the carcinogenicity of nitroso compounds, the relationship between processed meat and stomach cancer could be due to the high salt content of processed meats.

2.4. Salt intake and cancer

Early ecological studies showed a positive association between salt intake and stomach cancer, and this is consistent with the results of observational studies (Table 2). The evidence is consistent across studies investigating the intakes of salt, pickled vegetables, and other salty foods [4,100,101].

Animal and human studies have provided evidence of a synergistic effect of dietary salt intake and *Helicobacter pylori* infection on gastric carcinogenesis, and there is some evidence that salt intake may promote or enhance the carcinogenic effect of nitroso compounds [43].

2.5. Dietary supplements and cancer

Associations observed in epidemiological studies between dietary intake or biomarkers of exposure to certain micronutrients and cancer risk prompted several randomised controlled trials, most of which did not demonstrate a protective effect. Moreover, an unexpected increased cancer risk was observed in some trials.

Specifically, an increased risk of lung cancer was observed in men receiving beta-carotene supplements in the Alpha-Tocopherol, Beta-Carotene (ATBC) Cancer Prevention Study [102] and in the Beta-Carotene and Retinol Efficacy Trial (CARET) [103]. Beta-carotene was thought to reduce cancer risk by reducing oxidative damage, but it is now recognised that high doses of beta-carotene can result in pro-oxidant and anti-apoptotic effects, especially when cells are simultaneously subjected to tobacco smoke. Subsequent randomised controlled trials have confirmed the lack of a protective effect of beta-carotene supplementation on cancer risk [104]. An increased risk of prostate cancer was also observed in the follow-up of the Selenium and Vitamin E Cancer Prevention Trial (SELECT) in patients receiving high doses of vitamin E [105].

Total calcium intake is consistently associated with reduced risk of colorectal cancer in observational studies. Trials have shown a protective effect of calcium supplementation on colorectal adenomas, a precursor of colorectal cancer, but not against colorectal cancer itself [106]. Randomised controlled trials of folic acid supplementation with colorectal adenoma (reviewed by Ziegler [107]) or colorectal cancer [108] do not support the case that folic acid supplementation could prevent colorectal cancer. Several prospective studies suggest a trend towards a protective effect of folate (folic acid) intake on colon cancer risk [109], but in a recent meta-analysis of case-control studies nested in cohort studies, plasma or serum folate was not related to colorectal cancer risk [110]. It has been proposed that folate may play a dual role, with high concentrations promoting tumour development once premalignant lesions are established [111]. The evidence for an association between folate (folic acid) intake and increased or reduced cancer risk in humans is equivocal and does not support the recommendation of supplementation [112].

The null findings of dietary supplement interventions in randomised controlled trials may reflect lack of effect, or inappropriate type, timing, duration, or dose of the intervention.

The evidence for any benefit of vitamin and mineral supplementation for cancer prevention from observational studies is limited [113]. Dietary supplement use is not currently recommended for cancer prevention [4].

3. Justification for recommendation

3.1. Benefits for cancer prevention

There is evidence that diets characterised by high intake of plant foods (fruits, vegetables, pulses and whole-grain foods), low intake of red and processed meats, low intake of sugary foods, and avoidance of high salt intake are related to a lower risk of several cancers (Table 2). Recent data from the EPIC study show that compliance with a “healthy diet” such as recommended by the WCRF/AICR report (Table 1) is associated with a reduction in overall cancer risk (5% risk reduction for adherence to each additional recommendation), with the largest reductions for stomach, endometrial, oesophageal, colorectal, and mouth, pharynx, and larynx cancers (12–16% risk reductions) [114]. In two North American studies, compliance with the WCRF/AICR recommendations was associated with reductions in aggressive prostate cancer (13%) [115] and cancer mortality (10%) [116]. The Mediterranean dietary pattern is also consistently associated with significant reductions in cancer mortality (10%), and specifically with a lower risk of aerodigestive and colorectal cancers (reviewed by Schwingshackl and Hoffmann [117]).

3.2. Benefits on cancer prognosis

Recent reviews have shown that diet may modify biomarkers of cancer progression in individuals who have been treated for cancer [118,119]. The number of studies investigating the influence of diet after cancer diagnosis and cancer outcomes is limited. Current recommendations for cancer survivors are based on the recommendations to reduce cancer risk and emphasise achieving and maintaining a healthy weight, regular physical activity, and a diet rich in vegetables, fruits and whole grains, and limited in red meat and alcohol [4].

In observational studies, better pre- or post-diagnosis diet quality (in general dietary patterns rich in plant foods) has been found to be associated with reduced risk of death after diagnosis of breast cancer [120,121], colorectal cancer [122] and head and neck cancers [123]. A Western dietary pattern – characterised by high intakes of meat, fat, refined grains, and dessert – was associated with a higher risk of recurrence and mortality among patients with stage III colon cancer treated with surgery and adjuvant chemotherapy [124]. Overall, no benefit for cancer-related death has been observed in studies in cancer survivors, but such studies are difficult to interpret due to high risk of confounding or reverse causation.

Two dietary intervention studies among women diagnosed with breast cancer, the Women’s Healthy Eating and Living Trial (WHEL) and the Women’s Intervention Nutrition Study (WINS), found that dietary interventions among breast cancer survivors without weight loss or increase in physical activity do not improve breast cancer prognosis. WHEL focused on a plant-based dietary pattern that included a reduction in dietary fat, while WINS focused on reduced dietary fat intake. Secondary analyses in WHEL showed that the dietary intervention pattern was associated with a reduced risk of second breast cancer events among women with early-stage breast cancer who reported no hot flashes at baseline, and that higher vegetable intake was associated with reduced breast cancer recurrence in tamoxifen users [125,126].

3.3. Benefits for other diseases

Higher intakes of fruits, vegetables and fibre have been consistently associated with lower risk of coronary heart disease

Box 1. European Code Against Cancer.

EUROPEAN CODE AGAINST CANCER

12 ways to reduce your cancer risk

1. Do not smoke. Do not use any form of tobacco
2. Make your home smoke free. Support smoke-free policies in your workplace
3. Take action to be a healthy body weight
4. Be physically active in everyday life. Limit the time you spend sitting
5. Have a healthy diet:
 - Eat plenty of whole grains, pulses, vegetables and fruits
 - Limit high-calorie foods (foods high in sugar or fat) and avoid sugary drinks
 - Avoid processed meat; limit red meat and foods high in salt
6. If you drink alcohol of any type, limit your intake. Not drinking alcohol is better for cancer prevention
7. Avoid too much sun, especially for children. Use sun protection. Do not use sunbeds
8. In the workplace, protect yourself against cancer-causing substances by following health and safety instructions
9. Find out if you are exposed to radiation from naturally high radon levels in your home; take action to reduce high radon levels
10. For women:
 - Breastfeeding reduces the mother's cancer risk. If you can, breastfeed your baby
 - Hormone replacement therapy (HRT) increases the risk of certain cancers. Limit use of HRT
11. Ensure your children take part in vaccination programmes for:
 - Hepatitis B (for newborns)
 - Human papillomavirus (HPV) (for girls)
12. Take part in organised cancer screening programmes for:
 - Bowel cancer (men and women)
 - Breast cancer (women)
 - Cervical cancer (women)

The European Code Against Cancer focuses on actions that individual citizens can take to help prevent cancer. Successful cancer prevention requires these individual actions to be supported by governmental policies and actions.

in prospective studies (reviewed by Mente et al. [127]). A dietary pattern characterised by high intake of fruits and vegetables, and low intake of processed meat, sugar-sweetened beverages and refined cereal products, have been associated with lower risk of type 2 diabetes mellitus [128]. Numerous studies have shown that greater adherence to a Mediterranean diet is associated with a reduction in overall and cardiovascular disease mortality [129]. High salt intake increases blood pressure [130], which is a strong risk factor for cardiovascular and renal disease [131]. Processed meats have been found to be associated with higher incidence of coronary heart disease and diabetes mellitus [132].

In the EPIC study, participants with a lifestyle in agreement with the WCRF/AICR recommendations for cancer prevention had a 34% lower risk of death (95%CI: 0.59–0.75) compared with participants within the lowest agreement to the recommendations [133].

The 4th European Code against Cancer (Box 1) Nutrition Working Group has developed the recommendation of having a healthy diet in order to reduce the personal risk of getting cancer: “Have a healthy diet:

- eat plenty of whole grains, pulses, vegetables and fruits;
- limit high-calorie foods (foods high in sugar or fat) and avoid sugary drinks;
- avoid processed meat; limit red meat and foods high in salt.”

Conflict of interest

The authors declare no conflict of interest.

Authorship contribution

All the authors substantially contributed to the conception of the manuscript. Teresa Norat, Chiara Scoccianti and Marie-Christine Boutron-Ruault drafted the manuscript. All authors were involved in the revision of the manuscript.

Acknowledgments

The European Code Against Cancer project was co-funded by the European Union [grant agreement numbers: 2011 53 05; 2010 53 04 and 2007IARC01] and the International Agency for Research on Cancer. The authors alone are responsible for the views expressed in this paper.

References

- [1] Doll R, Peto R. The causes of cancer: quantitative estimates of avoidable risks of cancer in the United States today. *J Natl Cancer Inst* 1981;66(6):1191–308.
- [2] Armstrong B, Doll R. Environmental factors and cancer incidence and mortality in different countries, with special reference to dietary practices. *Int J Cancer* 1975;15(4):617–31.
- [3] Lippman SM, Hawk ET. Cancer prevention: from 1727 to milestones of the past 100 years. *Cancer Res* 2009;69(13):5269–84.
- [4] World Cancer Research Fund/American Institute for Cancer Research. Food, nutrition, physical activity and the prevention of cancer: a global perspective. Washington, DC: World Cancer Research Fund/American Institute for Cancer Research, 2007.
- [5] World Cancer Research Fund/American Institute for Cancer Research. Continuous update project; 2014. http://www.dietandcancerreport.org/cancer_resource_center/cup_summaries.php.
- [6] Wiseman M. The second World Cancer Research Fund/American Institute for Cancer Research expert report. Food, nutrition, physical activity, and the prevention of cancer: a global perspective. *Proc Nutr Soc* 2008;67(3):253–6.
- [7] World Cancer Research Fund/American Institute for Cancer Research. Cancer preventability estimates for food, nutrition, body fatness, and physical activity; 2014. http://www.wcrf.org/cancer_statistics/preventability_estimates/preventability_estimates_food.php.
- [8] Hursting SD, Lavigne JA, Berrigan D, Perkins SN, Barrett JC. Calorie restriction, aging, and cancer prevention: mechanisms of action and applicability to humans. *Annu Rev Med* 2003;54:131–52.
- [9] Anderson AS, Key TJ, Norat T, Scoccianti C, Cecchini M, Berrino F, Boutron-Ruault M-C, Espina C, Leitzmann M, Powers H, Wiseman M, Romieu I. European Code against Cancer 4th edition: Obesity, Body Fatness and Cancer. *Cancer Epidemiology* 2015;39:S34–45.
- [10] Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell* 2011;144(5):646–74.
- [11] Diet, nutrition, and the prevention of chronic diseases. Report of a WHO Study Group. *World Health Organ Tech Rep Ser* 1990;797:1–204.
- [12] Rolls BJ. The relationship between dietary energy density and energy intake. *Physiol Behav* 2009;97(5):609–15.
- [13] Micha R, Mozaffarian D. Trans fatty acids: effects on metabolic syndrome, heart disease and diabetes. *Nat Rev Endocrinol* 2009;5(6):335–44.
- [14] Chajes V, Thiebaut AC, Rotival M, Gauthier E, Maillard V, Boutron-Ruault MC, et al. Association between serum trans-monounsaturated fatty acids and breast cancer risk in the E3N-EPIC Study. *Am J Epidemiol* 2008;167(11):1312–20.
- [15] Wang J, John EM, Horn-Ross PL, Ingles SA. Dietary fat, cooking fat, and breast cancer risk in a multiethnic population. *Nutr Cancer* 2008;60(4):492–504.
- [16] Chavarro JE, Stampfer MJ, Campos H, Kurth T, Willett WC, Ma J. A prospective study of trans-fatty acid levels in blood and risk of prostate cancer. *Cancer Epidemiol Biomarkers Prev* 2008;17(11):95–101.
- [17] Chajes V, Jenab M, Romieu I, Ferrari P, Dahm CC, Overvad K, et al. Plasma phospholipid fatty acid concentrations and risk of gastric adenocarcinomas in the European Prospective Investigation into Cancer and Nutrition (EPIC-EURGAST). *Am J Clin Nutr* 2011;94(5):1304–13.
- [18] Vinikoor LC, Schroeder JC, Millikan RC, Satia JA, Martin CF, Ibrahim J, et al. Consumption of trans-fatty acid and its association with colorectal adenomas. *Am J Epidemiol* 2008;168(3):289–97.
- [19] World Health Organization. Global strategy on diet, physical activity and health: a framework to monitor and evaluate implementation. Geneva, Switzerland: World Health Organization, 2006.
- [20] Ceriello A, Bortolotti N, Motz E, Pieri C, Marra M, Tonutti L, et al. Meal-induced oxidative stress and low-density lipoprotein oxidation in diabetes: the possible role of hyperglycemia. *Metabolism* 1999;48(12):1503–8.
- [21] Hu FB. Resolved: there is sufficient scientific evidence that decreasing sugar-sweetened beverage consumption will reduce the prevalence of obesity and obesity-related diseases. *Obes Rev* 2013;14(8):606–19.
- [22] Clayton PE, Banerjee I, Murray PG, Renehan AG. Growth hormone, the insulin-like growth factor axis, insulin and cancer risk. *Nat Rev Endocrinol* 2011;7(1):11–24.
- [23] Cross AJ, Pollock JR, Bingham SA. Haem, not protein or inorganic iron, is responsible for endogenous intestinal N-nitrosation arising from red meat. *Cancer Res* 2003;63(10):2358–60.
- [24] Bastide NM, Pierre FH, Corpet DE. Heme iron from meat and risk of colorectal cancer: a meta-analysis and a review of the mechanisms involved. *Cancer Prev Res (Phila)* 2011;4(2):177–84.
- [25] Sugimura T, Wakabayashi K, Nakagama H, Nagao M. Heterocyclic amines: mutagens/carcinogens produced during cooking of meat and fish. *Cancer Sci* 2004;95(4):290–9.

- [26] IARC monographs on the evaluation of carcinogenic risks to humans. Ingested nitrate and nitrite, and cyanobacterial peptide toxins. IARC Monogr Eval Carcinog Risks Hum 2010; 94:v-vii, 1-412:v-412.
- [27] Gilsing AM, Berndt SI, Ruder EH, Graubard BI, Ferrucci LM, Burdett L, et al. Meat-related mutagen exposure, xenobiotic metabolizing gene polymorphisms and the risk of advanced colorectal adenoma and cancer. *Carcinogenesis* 2012;33(7):1332–9.
- [28] Brevik A, Joshi AD, Corral R, Onland-Moret NC, Siegmund KD, Le ML, et al. Polymorphisms in base excision repair genes as colorectal cancer risk factors and modifiers of the effect of diets high in red meat. *Cancer Epidemiol Biomarkers Prev* 2010;19(12):3167–73.
- [29] Andersen V, Ostergaard M, Christensen J, Overvad K, Tjønneland A, Vogel U. Polymorphisms in the xenobiotic transporter Multidrug Resistance 1 (MDR1) and interaction with meat intake in relation to risk of colorectal cancer in a Danish prospective case-cohort study. *BMC Cancer* 2009;9:407–9. <http://dx.doi.org/10.1186/1471-2407-9-407>.
- [30] Adlercreutz H, Hockerstedt K, Bannwart C, Bloigu S, Hamalainen E, Fotsis T, et al. Effect of dietary components, including lignans and phytoestrogens, on enterohepatic circulation and liver metabolism of estrogens and on sex hormone binding globulin (SHBG). *J Steroid Biochem* 1987;27(4–6):1135–44.
- [31] Monroe KR, Murphy SP, Henderson BE, Kolonel LN, Stanczyk FZ, Adlercreutz H, et al. Dietary fiber intake and endogenous serum hormone levels in naturally postmenopausal Mexican American women: the Multiethnic Cohort Study. *Nutr Cancer* 2007;58(2):127–35.
- [32] Maskarinec G, Morimoto Y, Takata Y, Murphy SP, Stanczyk FZ. Alcohol and dietary fibre intakes affect circulating sex hormones among premenopausal women. *Public Health Nutr* 2006;9(7):875–81.
- [33] Nangia-Makker P, Hogan V, Honjo Y, Baccharini S, Tait L, Bresalier R, et al. Inhibition of human cancer cell growth and metastasis in nude mice by oral intake of modified citrus pectin. *J Natl Cancer Inst* 2002;94(24):1854–62.
- [34] Mueller SO, Simon S, Chae K, Metzler M, Korach KS. Phytoestrogens and their human metabolites show distinct agonistic and antagonistic properties on estrogen receptor alpha (ERalpha) and ERbeta in human cells. *Toxicol Sci* 2004;80(1):14–25.
- [35] Penttinen P, Jaehrling J, Damdimopoulos AE, Inzunza J, Lemmen JG, van der Saag P, et al. Diet-derived polyphenol metabolite enterolactone is a tissue-specific estrogen receptor activator. *Endocrinology* 2007;148(10):4875–86.
- [36] Adlercreutz H. Phytoestrogens and breast cancer. *J Steroid Biochem Mol Biol* 2002;83(1–5):113–8.
- [37] Yoon JH, Baek SJ. Molecular targets of dietary polyphenols with anti-inflammatory properties. *Yonsei Med J* 2005;46(5):585–96.
- [38] Zhang Y. Cancer-preventive isothiocyanates: measurement of human exposure and mechanism of action. *Mutat Res* 2004;555(1–2):173–90.
- [39] Pathak SK, Sharma RA, Mellon JK. Chemoprevention of prostate cancer by diet-derived antioxidant agents and hormonal manipulation (Review). *Int J Oncol* 2003;22(1):5–13.
- [40] Dashwood RH, Myzak MC, Ho E. Dietary HDAC inhibitors: time to rethink weak ligands in cancer chemoprevention? *Carcinogenesis* 2006;27(2):344–9.
- [41] Chung MY, Lim TG, Lee KW. Molecular mechanisms of chemopreventive phytochemicals against gastroenterological cancer development. *World J Gastroenterol* 2013;19(7):984–93.
- [42] Beevers DG, Lip GY, Blann AD. Salt intake and *Helicobacter pylori* infection. *J Hypertens* 2004;22(8):1475–7.
- [43] D'Elia L, Galletti F, Strazzullo P. Dietary salt intake and risk of gastric cancer. *Cancer Treat Res* 2014;159:83–95. http://dx.doi.org/10.1007/978-3-642-38007-5_6.
- [44] Elmadafa I, Meyer A, Nowak V, Hasenegger V, Putz P, Verstraeten R, et al. European Nutrition and Health Report 2009. *Forum Nutr* 2009;62:1–405. 10.1159/000242367. Epub;2009 Sep 21.
- [45] Slimani N, Fahey M, Welch AA, Wirfalt E, Stripp C, Bergstrom E, et al. Diversity of dietary patterns observed in the European Prospective Investigation into Cancer and Nutrition (EPIC) project. *Public Health Nutr* 2002;5(6B):1311–28.
- [46] Naska A, Berg MA, Cuadrado C, Freisling H, Gedrich K, Gregoric M, et al. Food balance sheet and household budget survey dietary data and mortality patterns in Europe. *Br J Nutr* 2009;102(1):166–71.
- [47] Naska A, Bountziouka V, Trichopoulou A. Soft drinks: time trends and correlates in twenty-four European countries. A cross-national study using the DAFNE (Data Food Networking) databank. *Public Health Nutr* 2010;13(9):1346–55.
- [48] Riboli E, Norat T. Epidemiologic evidence of the protective effect of fruit and vegetables on cancer risk. *Am J Clin Nutr* 2003;78(3 Suppl.):559S–69S.
- [49] International Agency for Research on Cancer. Fruit and vegetables. Lyon: IARC Press, 2003.
- [50] Chuang SC, Jenab M, Heck JE, Bosetti C, Talamini R, Matsuo K, et al. Diet and the risk of head and neck cancer: a pooled analysis in the INHANCE consortium. *Cancer Causes Control* 2012;23(1):69–88.
- [51] Liu J, Wang J, Leng Y, Lv C. Intake of fruit and vegetables and risk of esophageal squamous cell carcinoma: a meta-analysis of observational studies. *Int J Cancer* 2013;133(2):473–85.
- [52] Iso H, Kubota Y. Nutrition and disease in the Japan Collaborative Cohort Study for Evaluation of Cancer (JACC). *Asian Pac J Cancer Prev* 2007;8(Suppl.):35–80.
- [53] Fan Y, Yuan JM, Wang R, Gao YT, Yu MC. Alcohol, tobacco, and diet in relation to esophageal cancer: the Shanghai Cohort Study. *Nutr Cancer* 2008;60(3):354–63.
- [54] Steevens J, Schouten LJ, Goldbohm RA, van den Brandt PA. Vegetables and fruits consumption and risk of esophageal and gastric cancer subtypes in the Netherlands Cohort Study. *Int J Cancer* 2011;129(11):2681–93.
- [55] Gonzalez CA, Pera G, Agudo A, et al. Fruit and vegetable intake and the risk of stomach and oesophagus adenocarcinoma in the European Prospective Investigation into Cancer and Nutrition (EPIC-EURGAST). *Int J Cancer* 2006;118(10):2559–66.
- [56] Wang Q, Chen Y, Wang X, Gong G, Li G, Li C. Consumption of fruit, but not vegetables, may reduce risk of gastric cancer: results from a meta-analysis of cohort studies. *Eur J Cancer* 2014;14:10.
- [57] Bae JM, Lee EJ, Guyatt G. Citrus fruit intake and stomach cancer risk: a quantitative systematic review. *Gastric Cancer* 2008;11(1):23–32.
- [58] Gonzalez CA, Lujan-Barroso L, Bueno-de-Mesquita HB, Jenab M, Duell EJ, Agudo A, et al. Fruit and vegetable intake and the risk of gastric adenocarcinoma: a reanalysis of the European Prospective Investigation into Cancer and Nutrition (EPIC-EURGAST) study after a longer follow-up. *Int J Cancer* 2012;131(12):2910–9.
- [59] Epplein M, Shu XO, Xiang YB, Chow WH, Yang G, Li HL, et al. Fruit and vegetable consumption and risk of distal gastric cancer in the Shanghai Women's and Men's Health studies. *Am J Epidemiol* 2010;172(4):397–406.
- [60] Jenab M, Riboli E, Ferrari P, Sabate J, Slimani N, Norat T, et al. Plasma and dietary vitamin C levels and risk of gastric cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC-EURGAST). *Carcinogenesis* 2006;27(11):2250–7.
- [61] Lam TK, Freedman ND, Fan JH, Qiao YL, Dawsey SM, Taylor PR, et al. Pre-diagnostic plasma vitamin C and risk of gastric adenocarcinoma and esophageal squamous cell carcinoma in a Chinese population. *Am J Clin Nutr* 2013;98(5):1289–97.
- [62] Freedman ND, Subar AF, Hollenbeck AR, Leitzmann MF, Schatzkin A, Abnet CC. Fruit and vegetable intake and gastric cancer risk in a large United States prospective cohort study. *Cancer Causes Control* 2008;19(5):459–67.
- [63] Zhou Y, Zhuang W, Hu W, Liu GJ, Wu TX, Wu XT. Consumption of large amounts of Allium vegetables reduces risk for gastric cancer in a meta-analysis. *Gastroenterology* 2011;141(1):80–9.
- [64] Ma JL, Zhang L, Brown LM, Li JY, Shen L, Pan KF, et al. Fifteen-year effects of *Helicobacter pylori*, garlic, and vitamin treatments on gastric cancer incidence and mortality. *J Natl Cancer Inst* 2012;104(6):488–92.
- [65] Buchner FL, Bueno-de-Mesquita HB, Ros MM, Overvad K, Dahm CC, Hansen L, et al. Variety in fruit and vegetable consumption and the risk of lung cancer in the European prospective investigation into cancer and nutrition. *Cancer Epidemiol Biomarkers Prev* 2010;19(9):2278–86.
- [66] Wright ME, Park Y, Subar AF, Freedman ND, Albanes D, Hollenbeck A, et al. Intakes of fruit, vegetables, and specific botanical groups in relation to lung cancer risk in the NIH-AARP Diet and Health Study. *Am J Epidemiol* 2008;168(9):1024–34.
- [67] Gallicchio L, Boyd K, Matanoski G, Tao XG, Chen L, Lam TK, et al. Carotenoids and the risk of developing lung cancer: a systematic review. *Am J Clin Nutr* 2008;88(2):372–83.
- [68] Takata Y, Xiang YB, Yang G, Li H, Gao J, Cai H, et al. Intakes of fruits, vegetables, and related vitamins and lung cancer risk: results from the Shanghai Men's Health Study (2002–2009). *Nutr Cancer* 2013;65(1):51–61.
- [69] Epplein M, Franke AA, Cooney RV, Morris JS, Wilkens LR, Goodman MT, et al. Association of plasma micronutrient levels and urinary isoprostane with risk of lung cancer: the multiethnic cohort study. *Cancer Epidemiol Biomarkers Prev* 2009;18(7):1962–70.
- [70] Aune D, Lau R, Chan DS, Vieira R, Greenwood DC, Kampman E, et al. Nonlinear reduction in risk for colorectal cancer by fruit and vegetable intake based on meta-analysis of prospective studies. *Gastroenterology* 2011;141(1):106–18.
- [71] Ruder EH, Thiebaut AC, Thompson FE, Potoschman N, Subar AF, Park Y, et al. Adolescent and mid-life diet: risk of colorectal cancer in the NIH-AARP Diet and Health Study. *Am J Clin Nutr* 2011;94(6):1607–19.
- [72] Aune D, Chan DS, Lau R, Vieira R, Greenwood DC, Kampman E, et al. Dietary fibre, whole grains, and risk of colorectal cancer: systematic review and dose-response meta-analysis of prospective studies. *BMJ* 2011;343:d6617. <http://dx.doi.org/10.1136/bmj.d6617>.
- [73] Curran J. The nutritional value and health benefits of pulses in relation to obesity, diabetes, heart disease and cancer. *Br J Nutr* 2012;108(Suppl. 1):S1–2. <http://dx.doi.org/10.1017/S0007114512003534>.
- [74] Aune D, Chan DS, Greenwood DC, et al. Dietary fiber and breast cancer risk: a systematic review and meta-analysis of prospective studies. *Ann Oncol* 2012;23(6):1394–402.
- [75] Coleman HG, Murray LJ, Hicks B, Bhat SK, Kubo A, Corley DA, et al. Dietary fiber and the risk of precancerous lesions and cancer of the esophagus: a systematic review and meta-analysis. *Nutr Rev* 2013;71(7):474–82.
- [76] Zhang Z, Xu G, Ma M, Yang J, Liu X. Dietary fiber intake reduces risk for gastric cancer: a meta-analysis. *Gastroenterology* 2013;145(1):113–20.
- [77] Liu L, Zhuang W, Wang RQ, Mukherjee R, Xiao SM, Chen Z, et al. Is dietary fat associated with the risk of colorectal cancer? A meta-analysis of 13 prospective cohort studies. *Eur J Nutr* 2011;50(3):173–84.
- [78] Prentice RL, Pettinger M, Tinker LF, Huang Y, Thomson CA, Johnson KC, et al. Regression calibration in nutritional epidemiology: example of fat density and total energy in relationship to postmenopausal breast cancer. *Am J Epidemiol* 2013;178(11):1663–72.

- [79] Bingham SA, Luben R, Welch A, Wareham N, Khaw KT, Day N. Are imprecise methods obscuring a relation between fat and breast cancer? *Lancet* 2003;362(9379):212–4.
- [80] Key TJ, Appleby PN, Cairns BJ, Luben R, Dahm CC, Akbaraly T, et al. Dietary fat and breast cancer: comparison of results from food diaries and food-frequency questionnaires in the UK Dietary Cohort Consortium. *Am J Clin Nutr* 2011;94(4):1043–52.
- [81] Turner LB. A meta-analysis of fat intake, reproduction, and breast cancer risk: an evolutionary perspective. *Am J Hum Biol* 2011;23(5):601–8.
- [82] Sieri S, Chiodini P, Agnoli C, Pala V, Berrino F, Trichopoulou A, et al. Dietary fat intake and development of specific breast cancer subtypes. *J Natl Cancer Inst* 2014;88(5):1304–12.
- [83] Prentice RL, Thomson CA, Caan B, Hubbell FA, Anderson GL, Beresford SA, et al. Low-fat dietary pattern and cancer incidence in the Women's Health Initiative Dietary Modification Randomized Controlled Trial. *J Natl Cancer Inst* 2007;99(20):1534–43.
- [84] Tasevska N, Park Y, Jiao L, Hollenbeck A, Subar AF, Potosichman N. Sugars and risk of mortality in the NIH-AARP diet and health study. *Am J Clin Nutr* 2014;99(5):1077–88.
- [85] InterAct Consortium, Romaguera D, Norat T, Wark PA, Vergnaud AC, Schulze MB, van Woudenberg GJ, et al. Consumption of sweet beverages and type 2 diabetes incidence in European adults: results from EPIC-InterAct. *Diabetologia* 2013;56(7):1520–30.
- [86] Aune D, Greenwood DC, Chan DS, Vieira R, Vieira AR, Navarro Rosenblatt DA, et al. Body mass index, abdominal fatness and pancreatic cancer risk: a systematic review and non-linear dose–response meta-analysis of prospective studies. *Ann Oncol* 2012;23(4):843–52.
- [87] Elena JW, Steplowski E, Yu K, Hartge P, Tobias GS, Brotzman MJ, et al. Diabetes and risk of pancreatic cancer: a pooled analysis from the pancreatic cancer cohort consortium. *Cancer Causes Control* 2013;24(1):13–25.
- [88] Gallus S, Turati F, Tavani A, Polesel J, Talamini R, Franceschi S, et al. Soft drinks, sweetened beverages and risk of pancreatic cancer. *Cancer Causes Control* 2011;22(1):33–9.
- [89] Genkinger JM, Li R, Spiegelman D, Anderson KE, Albanes D, Bergkvist L, et al. Coffee, tea, and sugar-sweetened carbonated soft drink intake and pancreatic cancer risk: a pooled analysis of 14 cohort studies. *Cancer Epidemiol Biomarkers Prev* 2012;21(2):305–18.
- [90] Aune D, Chan DS, Vieira AR, Navarro Rosenblatt DA, Vieira R, Greenwood DC, et al. Dietary fructose, carbohydrates, glycemic indices and pancreatic cancer risk: a systematic review and meta-analysis of cohort studies. *Ann Oncol* 2012;23(10):2536–46.
- [91] Vos MB, Kimmons JE, Gillespie C, Welsh J, Blanck HM. Dietary fructose consumption among US children and adults: the Third National Health and Nutrition Examination Survey. *Medscape J Med* 2008;10(7):160.
- [92] Zhang X, Albanes D, Beeson WL, van den Brandt PA, Buring JE, Flood A, et al. Risk of colon cancer and coffee, tea, and sugar-sweetened soft drink intake: pooled analysis of prospective cohort studies. *J Natl Cancer Inst* 2010;102(11):771–83.
- [93] Joosen AM, Kuhnle GG, Aspinall SM, Barrow TM, Lecommandeur E, Azqueta A, et al. Effect of processed and red meat on endogenous nitrosation and DNA damage. *Carcinogenesis* 2009;30(8):1402–7.
- [94] Choi Y, Giovannucci E, Lee JE. Glycaemic index and glycaemic load in relation to risk of diabetes-related cancers: a meta-analysis. *Br J Nutr* 2012;108(11):1934–47.
- [95] Chan DS, Lau R, Aune D, Vieira R, Greenwood DC, Kampman E, et al. Red and processed meat and colorectal cancer incidence: meta-analysis of prospective studies. *PLoS ONE* 2011;6(6):e20456.
- [96] Zhu H, Yang X, Zhang C, Zhu C, Tao G, Zhao L, et al. Red and processed meat intake is associated with higher gastric cancer risk: a meta-analysis of epidemiological observational studies. *PLoS ONE* 2013;8(8):e70955.
- [97] Larsson SC, Wolk A. Red and processed meat consumption and risk of pancreatic cancer: meta-analysis of prospective studies. *Br J Cancer* 2012;106(3):603–7.
- [98] Rohrmann S, Overvad K, Bueno-de-Mesquita HB, Jakobsen MU, Egeberg R, Tjønneland A, et al. Meat consumption and mortality – results from the European Prospective Investigation into Cancer and Nutrition. *BMC Med* 2013;11:63. <http://dx.doi.org/10.1186/1741-7015-11-63>.
- [99] Pan A, Sun Q, Bernstein AM, Schulze MB, Manson JE, Stampfer MJ, et al. Red meat consumption and mortality: results from 2 prospective cohort studies. *Arch Intern Med* 2012;172(7):555–63.
- [100] Ge S, Feng X, Shen L, Wei Z, Zhu Q, Sun J. Association between habitual dietary salt intake and risk of gastric cancer: a systematic review of observational studies. *Gastroenterol Res Pract* 2012;808120. <http://dx.doi.org/10.1155/2012/808120>. Epub; 2012 Oct 22. :808120.
- [101] D'Elia L, Rossi G, Ippolito R, Cappuccio FP, Strazzullo P. Habitual salt intake and risk of gastric cancer: a meta-analysis of prospective studies. *Clin Nutr* 2012;31(4):489–98.
- [102] Albanes D, Heinonen OP, Huttunen JK, Taylor PR, Virtamo J, Edwards BK, et al. Effects of alpha-tocopherol and beta-carotene supplements on cancer incidence in the Alpha-Tocopherol Beta-Carotene Cancer Prevention Study. *Am J Clin Nutr* 1995;62(6 Suppl.):1427S–30S.
- [103] Omenn GS, Goodman GE, Thornquist MD, Balmes J, Cullen MR, Glass A, et al. Risk factors for lung cancer and for intervention effects in CARET, the beta-carotene and retinol efficacy trial. *J Natl Cancer Inst* 1996;88(21):1550–9.
- [104] Druesne-Pecollo N, Latino-Martel P, Norat T, Barrandon E, Bertrais S, Galan P, et al. Beta-carotene supplementation and cancer risk: a systematic review and meta-analysis of randomized controlled trials. *Int J Cancer* 2010;127(1):172–84.
- [105] Klein EA, Thompson Jr IM, Tangen CM, Crowley JJ, Lucia MS, Goodman PJ, et al. Vitamin E and the risk of prostate cancer: the Selenium and Vitamin E Cancer Prevention Trial (SELECT). *JAMA* 2011;306(14):1549–56.
- [106] Keum N, Aune D, Greenwood DC, Ju W, Giovannucci EL. Calcium intake and colorectal cancer risk: dose–response meta-analysis of prospective observational studies. *Int J Cancer* 2014;135(8):1940–8.
- [107] Ziegler RG. Following up folate and its function in colorectal carcinogenesis. *J Natl Cancer Inst* 2012;104(20):1525–7.
- [108] Vollset SE, Clarke R, Lewington S, Ebbing M, Halsey J, Lonn E, et al. Effects of folic acid supplementation on overall and site-specific cancer incidence during the randomised trials: meta-analyses of data on 50,000 individuals. *Lancet* 2013;381(9871):1029–36.
- [109] Kim DH, Smith-Warner SA, Spiegelman D, Yaun SS, Colditz GA, Freudenheim JL, et al. Pooled analyses of 13 prospective cohort studies on folate intake and colon cancer. *Cancer Causes Control* 2010;21(11):1919–30.
- [110] Chuang SC, Rota M, Gunter MJ, Zeleniuch-Jacquotte A, Eussen SJ, Vollset SE, et al. Quantifying the dose–response relationship between circulating folate concentrations and colorectal cancer in cohort studies: a meta-analysis based on a flexible meta-regression model. *Am J Epidemiol* 2013;178(7):1028–37.
- [111] Kim YI. Role of folate in colon cancer development and progression. *J Nutr* 2003;133(11 Suppl. 1):3731S–9S.
- [112] The Scientific Advisory Committee on Nutrition Folate and Disease Prevention Report, 2006; <https://www.gov.uk/government/publications/sacn-folate-and-disease-prevention-report>
- [113] Fortmann SP, Burda BU, Senger CA, Lin JS, Whitlock EP. Vitamin and mineral supplements in the primary prevention of cardiovascular disease and cancer: an updated systematic evidence review for the U.S. Preventive Services Task Force. *Ann Intern Med* 2013;159(12):824–34.
- [114] Romaguera D, Vergnaud AC, Peeters PH, van Gils CH, Chan DS, Ferrari P, et al. Is concordance with World Cancer Research Fund/American Institute for Cancer Research guidelines for cancer prevention related to subsequent risk of cancer? Results from the EPIC study. *Am J Clin Nutr* 2012;96(1):150–63.
- [115] Arab L, Su J, Steck SE, Ang A, Fontham ET, Bensen JT, et al. Adherence to World Cancer Research Fund/American Institute for Cancer Research lifestyle recommendations reduces prostate cancer aggressiveness among African and Caucasian Americans. *Nutr Cancer* 2013;65(5):633–43.
- [116] Hastert TA, Beresford SA, Sheppard L, White E. Adherence to the WCRF/AICR cancer prevention recommendations and cancer-specific mortality: results from the Vitamins and Lifestyle (VITAL) Study. *Cancer Causes Control* 2014;25(5):541–52.
- [117] Schwingshackl L, Hoffmann G. Adherence to Mediterranean diet and risk of cancer: A systematic review and meta-analysis of observational studies. *Int J Cancer* 2014.
- [118] Jones LW, Demark-Wahnefried W. Diet, exercise, and complementary therapies after primary treatment for cancer. *Lancet Oncol* 2006;7(12):1017–26.
- [119] Pekmezci DW, Demark-Wahnefried W. Updated evidence in support of diet and exercise interventions in cancer survivors. *Acta Oncol* 2011;50(2):167–78.
- [120] George SM, Ballard-Barbash R, Shikany JM, Caan BJ, Freudenheim JL, Kroenke CH, et al. Better postdiagnosis diet quality is associated with reduced risk of death among postmenopausal women with invasive breast cancer in the women's health initiative. *Cancer Epidemiol Biomarkers Prev* 2014;23(4):575–83. <http://dx.doi.org/10.1158/1055-9965>.
- [121] Vrieling A, Buck K, Seibold P, Heinz J, Obi N, Flesch-Janys D, et al. Dietary patterns and survival in German postmenopausal breast cancer survivors. *Br J Cancer* 2013;108(1):188–92.
- [122] Pelser C, Arem H, Pfeiffer RM, Elena JW, Alfano CM, Hollenbeck AR, et al. Prediagnostic lifestyle factors and survival after colon and rectal cancer diagnosis in the National Institutes of Health (NIH)-AARP Diet and Health Study. *Cancer* 2014;120(10):1540–7. <http://dx.doi.org/10.1002/cncr.28573>.
- [123] Arthur AE, Peterson KE, Rozek LS, Taylor JM, Light E, Chepeha DB, et al. Pretreatment dietary patterns, weight status, and head and neck squamous cell carcinoma prognosis. *Am J Clin Nutr* 2013;97(2):360–8.
- [124] Meyerhardt JA, Niedzwiecki D, Hollis D, Saltz LB, Hu FB, Mayer RJ, et al. Association of dietary patterns with cancer recurrence and survival in patients with stage III colon cancer. *JAMA* 2007;298(7):754–64.
- [125] Gold EB, Pierce JP, Natarajan L, Stefanick ML, Laughlin GA, Caan BJ, et al. Dietary pattern influences breast cancer prognosis in women without hot flashes: the women's healthy eating and living trial. *J Clin Oncol* 2009;27(3):352–9.
- [126] Thomson CA, Rock CL, Thompson PA, Caan BJ, Cussler E, Flatt SW, et al. Vegetable intake is associated with reduced breast cancer recurrence in tamoxifen users: a secondary analysis from the Women's Healthy Eating and Living Study. *Breast Cancer Res Treat* 2011;125(2):519–27.
- [127] Mente A, de KL, Shannon HS, Anand SS. A systematic review of the evidence supporting a causal link between dietary factors and coronary heart disease. *Arch Intern Med* 2009;169(7):659–69.
- [128] Adherence to predefined dietary patterns and incident type 2 diabetes in European populations: EPIC-InterAct Study. *Diabetologia* 2014;57(2):321–33.
- [129] Sofi F, Cesari F, Abbate R, Gensini GF, Casini A. Adherence to Mediterranean diet and health status: meta-analysis. *BMJ* 2008;337:a1344. <http://dx.doi.org/10.1136/bmj.a1344>.

- [130] Appel LJ, Brands MW, Daniels SR, Karanja N, Elmer PJ, Sacks FM. Dietary approaches to prevent and treat hypertension: a scientific statement from the American Heart Association. *Hypertension* 2006;47(2):296–308.
- [131] Aaron KJ, Sanders PW. Role of dietary salt and potassium intake in cardiovascular health and disease: a review of the evidence. *Mayo Clin Proc* 2013;88(9):987–95.
- [132] Micha R, Wallace SK, Mozaffarian D. Red and processed meat consumption and risk of incident coronary heart disease, stroke, and diabetes mellitus: a systematic review and meta-analysis. *Circulation* 2010;121(21):2271–83.
- [133] Vergnaud AC, Romaguera D, Peeters PH, van Gils CH, Chan DS, Romieu J, et al. Adherence to the World Cancer Research Fund/American Institute for Cancer Research guidelines and risk of death in Europe: results from the European Prospective Investigation into Nutrition and Cancer cohort study1,4. *Am J Clin Nutr* 2013;97(5):1107–20.
- [134] Kim HJ, Lim SY, Lee JS, Park S, Shin A, Choi BY, et al. Fresh and pickled vegetable consumption and gastric cancer in Japanese and Korean populations: a meta-analysis of observational studies. *Cancer Sci* 2010;101(2):508–16.
- [135] Meng S, Zhang X, Giovannucci EL, Ma J, Fuchs CS, Cho E. No association between garlic intake and risk of colorectal cancer. *Cancer Epidemiol* 2013;37(2):152–5.
- [136] McCullough ML, Jacobs EJ, Shah R, Campbell PT, Gapstur SM. Garlic consumption and colorectal cancer risk in the CPS-II Nutrition Cohort. *Cancer Causes Control* 2012;23(10):1643–51.
- [137] Satia JA, Littman A, Slatore CG, Galanko JA, White E. Associations of herbal and specialty supplements with lung and colorectal cancer risk in the VITamins and Lifestyle study. *Cancer Epidemiol Biomarkers Prev* 2009;18(5):1419–28.
- [138] Lucenteforte E, Garavello W, Bosetti C, La VC. Dietary factors and oral and pharyngeal cancer risk. *Oral Oncol* 2009;45(6):461–7.
- [139] Soerjomataram I, Oomen D, Lemmens V, Oenema A, Benetou V, Trichopoulou A, et al. Increased consumption of fruit and vegetables and future cancer incidence in selected European countries. *Eur J Cancer* 2010;46(14):2563–80.
- [140] Ren JS, Kamangar F, Forman D, Islami F. Pickled food and risk of gastric cancer – a systematic review and meta-analysis of English and Chinese literature. *Cancer Epidemiol Biomarkers Prev* 2012;21(6):905–15.