

MTH 102 Calculus -II

Dr. Gizem SEYHAN ÖZTEPE

Topic-13-Chain Rule –Implicit Differentiation

Maximum and Minimum Problems



Chain Rule

Theorem

Let the function $z = f(x, y)$ has continuous partial derivatives f_x and f_y . If the functions $x = g(u, v)$ and $y = h(u, v)$ have partial derivatives with respect to u and v , then the function $z = f(g(u, v), h(u, v))$ has partial derivatives with respect to u and v

$$\frac{\partial f}{\partial u} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u}$$

$$\frac{\partial f}{\partial v} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial v}$$

Implicit Differentiation

Theorem:

Let the function $z=f(x,y)$ given by $F(x, y, z) = 0$. If the partial derivatives F_x and F_y are continuous and $F_z \neq 0$, then from the chain rule we obtain that

$$\frac{\partial F}{\partial x} \frac{dx}{dx} + \frac{\partial F}{\partial y} \frac{\partial y}{\partial x} + \frac{\partial F}{\partial z} \frac{\partial z}{\partial x} = 0$$

Since $\frac{dx}{dx}=1$ and $\frac{\partial y}{\partial x}=0$

$$\frac{\partial F}{\partial x} + \frac{\partial F}{\partial z} \frac{\partial z}{\partial x} = 0$$

So,

$$\frac{\partial z}{\partial x} = \frac{-F_x}{F_z}$$


Similarly taking the derivative with respect to y , we obtain

$$\frac{\partial z}{\partial y} = \frac{-F_y}{F_z}$$

We can summarize our results as follows

$$\frac{\partial z}{\partial x} = -\frac{\frac{\partial F}{\partial x}}{\frac{\partial F}{\partial z}} \quad \frac{\partial z}{\partial y} = -\frac{\frac{\partial F}{\partial y}}{\frac{\partial F}{\partial z}}$$

Maximum and Minimum Problems



Look at the hills and valleys in the graph of shown in Figure. There are two points (a, b) where f has a *local maximum*, that is, where $f(a, b)$ is larger than nearby values of $f(x, y)$. The larger of these two values is the *absolute maximum*. Likewise, f has two *local minima*, where $f(a, b)$ is smaller than nearby values. The smaller of these two values is the *absolute minimum*.

1 Definition A function of two variables has a **local maximum** at (a, b) if $f(x, y) \leq f(a, b)$ when (x, y) is near (a, b) . [This means that $f(x, y) \leq f(a, b)$ for all points (x, y) in some disk with center (a, b) .] The number $f(a, b)$ is called a **local maximum value**. If $f(x, y) \geq f(a, b)$ when (x, y) is near (a, b) , then f has a **local minimum** at (a, b) and $f(a, b)$ is a **local minimum value**.

- If the inequalities in Definition 1 hold for all points (x, y) in the domain of f , then f has an absolute maximum (or absolute minimum) at (a, b) .

2 Theorem If f has a local maximum or minimum at (a, b) and the first-order partial derivatives of f exist there, then $f_x(a, b) = 0$ and $f_y(a, b) = 0$.

Definition (Critical Point)

A point (a, b) is called a critical point of f if $f_x(a, b) = 0$ and $f_y(a, b) = 0$

We need to be able to determine whether or not a function has an extreme value (local min. or max.) at a critical point. The following test is given for this:

3 Second Derivatives Test Suppose the second partial derivatives of f are continuous on a disk with center (a, b) , and suppose that $f_x(a, b) = 0$ and $f_y(a, b) = 0$ [that is, (a, b) is a critical point of f]. Let

$$D = D(a, b) = f_{xx}(a, b) f_{yy}(a, b) - [f_{xy}(a, b)]^2$$

- (a) If $D > 0$ and $f_{xx}(a, b) > 0$, then $f(a, b)$ is a local minimum.
- (b) If $D > 0$ and $f_{xx}(a, b) < 0$, then $f(a, b)$ is a local maximum.
- (c) If $D < 0$, then $f(a, b)$ is not a local maximum or minimum.

NOTE 1 In case (c) the point (a, b) is called a **saddle point** of f and the graph of f crosses its tangent plane at (a, b) .

NOTE 2 If $D = 0$, the test gives no information: f could have a local maximum or local minimum at (a, b) , or (a, b) could be a saddle point of f .

NOTE 3 To remember the formula for D , it's helpful to write it as a determinant:

$$D = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{vmatrix} = f_{xx} f_{yy} - (f_{xy})^2$$