



# Calculus Lecture 1

Oktay Ölmez, Murat Şahin and Serhan Varma

# A brief summary of the concept of functions




# A brief summary of the concept of functions




## Definition

*A function is a rule that assigns to each element in a set  $A$  one and only one element in a set  $B$ .*

# A brief summary of the concept of functions




# A brief summary of the concept of functions




- The set  $A$  is called the domain of the function.

# A brief summary of the concept of functions



- The set  $A$  is called the domain of the function.
- If  $x$  is an element in the domain of a function  $f$ , then the element in  $B$  that  $f$  associates with  $x$  is written  $f(x)$  (read  $f$  of  $x$ ) and is called the value of  $f$  at  $x$ .

# A brief summary of the concept of functions



- The set  $A$  is called the domain of the function.
- If  $x$  is an element in the domain of a function  $f$ , then the element in  $B$  that  $f$  associates with  $x$  is written  $f(x)$  (read  $f$  of  $x$ ) and is called the value of  $f$  at  $x$ .
- The set comprising all the values assumed by  $y = f(x)$  as  $x$  takes on all possible values in its domain is called the range of the function  $f$ .

## Example

*An open box is to be made from a rectangular piece of cardboard 16 inches long and 10 inches wide by cutting away identical squares ( $x$  inches by  $x$  inches) from each corner and folding up the resulting flaps. Find an expression that gives the volume  $V$  of the box as a function of  $x$ . What is the domain of the function?*



## Example

*An open box is to be made from a rectangular piece of cardboard 16 inches long and 10 inches wide by cutting away identical squares ( $x$  inches by  $x$  inches) from each corner and folding up the resulting flaps. Find an expression that gives the volume  $V$  of the box as a function of  $x$ . What is the domain of the function?*

## Solution

$$V(x) = (16 - 2x)(10 - 2x)x$$

## Example

*An open box is to be made from a rectangular piece of cardboard 16 inches long and 10 inches wide by cutting away identical squares ( $x$  inches by  $x$  inches) from each corner and folding up the resulting flaps. Find an expression that gives the volume  $V$  of the box as a function of  $x$ . What is the domain of the function?*

## Solution

$$V(x) = (16 - 2x)(10 - 2x)x$$

*To find the domain, we need to consider the following inequalities:*

$$16 - 2x > 0 \quad 10 - 2x > 0 \quad x > 0$$

*Thus the domain is  $(0, 5)$ .*

## Example

*Find the domain of the each function:*

- (a)  $\sqrt{x-1}$
- (b)  $\frac{1}{x^2-4}$
- (c)  $x^2 + 3$

## Example

Find the domain of the each function:

- (a)  $\sqrt{x-1}$
- (b)  $\frac{1}{x^2-4}$
- (c)  $x^2 + 3$

## Answers

- (a)  $x \geq 1$
- (b)  $\mathbb{R} \setminus \{\pm 2\}$
- (c)  $\mathbb{R}$

# Graph of a Function of One Variable


## Definition

*The graph of a function  $f$  is the set of all points  $(x, y)$  in the  $xy$ -plane such that  $x$  is in the domain of  $f$  and  $y = f(x)$ .*

# Graph of a Function of One Variable

## Definition

*The graph of a function  $f$  is the set of all points  $(x, y)$  in the  $xy$ -plane such that  $x$  is in the domain of  $f$  and  $y = f(x)$ .*




## Example

*A Finance Company plans to open two branch offices 2 years from now in two separate locations: an industrial complex and a newly developed commercial center in the city. As a result of these expansion plans, The Company's total deposits during the next 5 years are expected to grow in accordance with the rule where*

$$f(x) = \begin{cases} \sqrt{2x} + 20 & \text{if } 0 \leq x \leq 2 \\ \frac{x^2}{2} + 20 & \text{if } 2 < x \leq 5 \end{cases}$$

*gives the total amount of money (in millions of dollars) on deposit with the company in year  $x$  ( $x = 0$  corresponds to the present). Sketch the graph of the function  $f$ .*

# Solution





# Some special functions: Absolute Value Function

## Definition (Absolute value function)

*The absolute value function is defined as*


$$f(x) = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

# Some special functions: Absolute Value Function

## Definition (Absolute value function)

*The absolute value function is defined as*

$$f(x) = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$




# Some special functions: Absolute Value Function

## Definition (Absolute value function)

*The absolute value function is defined as*

$$f(x) = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$



The domain of the absolute value function is the set of all real numbers and the range is the set of all positive real numbers including zero.

# Some special functions: Sign Function

## Definition (Sign function)

The sign or signum function,  $\text{sgn}$ , is defined according to


$$f(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -1 & \text{if } x < 0 \end{cases}$$

# Some special functions: Sign Function

## Definition (Sign function)

The sign or signum function,  $\text{sgn}$ , is defined according to

$$f(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -1 & \text{if } x < 0 \end{cases}$$



# Some special functions: Floor Function

## Definition (Floor function)

*Floor function, denoted by  $\lfloor x \rfloor$ , is defined as the greatest integer less than or equal to any real number. For example,  $\lfloor 2.3 \rfloor = 2$ ,  $\lfloor 0.4 \rfloor = 0$  and  $\lfloor -3.1 \rfloor = -4$ . This function has an infinite number of breaks or steps-one at each integer value in its domain.*

# Some special functions: Floor Function

## Definition (Floor function)

*Floor function, denoted by  $\lfloor x \rfloor$ , is defined as the greatest integer less than or equal to any real number. For example,  $\lfloor 2.3 \rfloor = 2$ ,  $\lfloor 0.4 \rfloor = 0$  and  $\lfloor -3.1 \rfloor = -4$ . This function has an infinite number of breaks or steps-one at each integer value in its domain.*

