

Calculus

Lecture 3

Oktay Ölmez, Murat Şahin and Serhan Varma

Intermediate Value Theorem

We say f is continuous on the interval $[a, b]$ if

- f is continuous at every point that belongs to the interval (a, b) .
- $\lim_{x \rightarrow a^+} f(x) = f(a)$
- $\lim_{x \rightarrow b^-} f(x) = f(b)$

Intermediate Value Theorem

We say f is continuous on the interval $[a, b]$ if

- f is continuous at every point that belongs to the interval (a, b) .
- $\lim_{x \rightarrow a^+} f(x) = f(a)$
- $\lim_{x \rightarrow b^-} f(x) = f(b)$

Theorem (IVT)

Let f be a continuous function on $[a, b]$ and W be a number between $f(a)$ and $f(b)$. Then, there is at least one c between a and b such that $f(c) = W$.


Example

Is there any real number c between 0 and $\pi/2$ such that $c = \cos(c)$?

IVT Example

Example

Is there any real number c between 0 and $\pi/2$ such that $c = \cos(c)$?



Example

Show that $p(x) = 2x^3 - 5x^2 - 10x + 5$ has a root somewhere between -1 and 2 .

Types of discontinuity

- In a **removable discontinuity** $\lim_{x \rightarrow c} f(x)$ exists, but $\lim_{x \rightarrow c} f(x) \neq f(c)$.
- In a **jump discontinuity**, the right-hand and left-hand limits both exist, but are not equal.
- An **infinite discontinuity** exists when one of the one-sided limits of the function is infinite.
- An **oscillating discontinuity** exists when the values of the function appear to be approaching two or more values simultaneously.

