

Calculus

Lecture 4

Oktay Ölmez, Murat Şahin and Serhan Varma


Meaning





- imitative of the work of another artist, writer, etc., and usually disapproved of for that reason.
- (of a product) having a value deriving from an underlying variable asset.
- something which is based on another source.

why derivative?



Slope of tangent line

Consider the graph of $y = f(x)$.

Slope of tangent line

Consider the graph of $y = f(x)$.

- $m_{sec} = \frac{f(x+h) - f(x)}{h}$.

- $m_{tan} = \lim_{h \rightarrow 0} m_{sec} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ (provided that this limit exists and not equals to either ∞ or $-\infty$).

Slope of tangent line

Consider the graph of $y = f(x)$.

- $m_{sec} = \frac{f(x+h) - f(x)}{h}$.
- $m_{tan} = \lim_{h \rightarrow 0} m_{sec} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ (provided that this limit exists and not equals to either ∞ or $-\infty$).

Suppose $f(x)$ is the position function of a moving object.

- $v_{avg} = \frac{f(x+h) - f(x)}{h}$.
- $v = \lim_{h \rightarrow 0} v_{avg} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$. (v is the instantaneous velocity)

Secant to Tangent

Definition

Derivative of a function f is another function f' whose value at number x is given by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Example

Example

Apply the definition to find the derivative of the function $f(x) = x^2$.

Solution

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\&= \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h} \\&= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - x^2}{h} \\&= \lim_{h \rightarrow 0} \frac{h(2x+h)}{h} \\&= \lim_{h \rightarrow 0} 2x + h = 2x.\end{aligned}$$

Differentiability implies continuity

Theorem

If $f'(c)$ exists then f is continuous at c .

Differentiability implies continuity

Theorem

If $f'(c)$ exists then f is continuous at c .


But the converse may not be true !

Differentiability implies continuity

Theorem

If $f'(c)$ exists then f is continuous at c .

But the converse may not be true !



$f(x) = |x|$ is not differentiable at $x = 0$

By definition, we need to show the following limit exists at $x = 0$.

$$f'(x) = \lim_{h \rightarrow 0} \frac{|x + h| - |x|}{h}$$

Thus we just need to consider,

$$\lim_{h \rightarrow 0} \frac{|0 + h| - |0|}{h} = \lim_{h \rightarrow 0} \frac{|h|}{h}.$$

But, it is evident that this limit DNE.

Example

The rate of change of electric charge with respect to time is called current. Suppose that $\frac{t^3}{3} + t$ coulombs of charge flow through a wire in t seconds. Find the current in amperes after 3 seconds. When will a 26-amperes fuse in the line?

Δx and Δy

Let $y = f(x)$. Then,

Δx and Δy

Let $y = f(x)$. Then,

- Δx means change in x .

Δx and Δy

Let $y = f(x)$. Then,

- Δx means change in x .
- If $x_1 = 2$ and $x_2 = 4$, Then $\Delta x = 4 - 2 = 2$.

Δx and Δy

Let $y = f(x)$. Then,

- Δx means change in x .
- If $x_1 = 2$ and $x_2 = 4$, Then $\Delta x = 4 - 2 = 2$.
- Δy means change in y .

Δx and Δy

Let $y = f(x)$. Then,

- Δx means change in x .
- If $x_1 = 2$ and $x_2 = 4$, Then $\Delta x = 4 - 2 = 2$.
- Δy means change in y .
- If $x_1 = 2$ and $x_2 = 4$, Then $\Delta y = f(4) - f(2)$.


Δx and Δy

Let $y = f(x)$. Then,

- Δx means change in x .
- If $x_1 = 2$ and $x_2 = 4$, Then $\Delta x = 4 - 2 = 2$.
- Δy means change in y .
- If $x_1 = 2$ and $x_2 = 4$, Then $\Delta y = f(4) - f(2)$.
- $m_{sec} = \frac{\Delta y}{\Delta x}$.
- $m_{tan} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} = \frac{dy}{dx}$

Change yourself!

Change yourself!



Equation of a Tangent Line

Equation of a Tangent Line

Example

Find the equation of the tangent line to the graph of $y = \frac{1}{x}$ at $x = 1$.