

Calculus

Lecture 6

Oktay Ölmez, Murat Şahin and Serhan Varma

Maxima and minima

Let S , the domain of f contain the point c .

- $f(c)$ is the maximum value of f on S if $f(c) \geq f(x)$ for all $x \in S$.

Maxima and minima

Let S , the domain of f contain the point c .

- $f(c)$ is the maximum value of f on S if $f(c) \geq f(x)$ for all $x \in S$.
- $f(c)$ is the minimum value of f on S if $f(c) \leq f(x)$ for all $x \in S$.

Maxima and minima

Let S , the domain of f contain the point c .

- $f(c)$ is the maximum value of f on S if $f(c) \geq f(x)$ for all $x \in S$.
- $f(c)$ is the minimum value of f on S if $f(c) \leq f(x)$ for all $x \in S$.
- $f(c)$ is an extreme value of f on S if it is either the maximum value or the minimum value.

Maxima and minima

Let S , the domain of f contain the point c .

- $f(c)$ is the maximum value of f on S if $f(c) \geq f(x)$ for all $x \in S$.
- $f(c)$ is the minimum value of f on S if $f(c) \leq f(x)$ for all $x \in S$.
- $f(c)$ is an extreme value of f on S if it is either the maximum value or the minimum value.
- the function we want to maximize or minimize is called objective function.

Does f have a maximum or a minimum value on S ?

If f is continuous on a closed interval $[a, b]$, then f attains both a maximum value and minimum value there.

Where do extreme values occur?

Let S , the domain of f , contains the point c .

- End points of the interval

Where do extreme values occur?

Let S , the domain of f , contains the point c .

- End points of the interval
- stationary points: If c is a point which $f'(c) = 0$, we call c a stationary point

Where do extreme values occur?

Let S , the domain of f , contains the point c .

- End points of the interval
- stationary points: If c is a point which $f'(c) = 0$, we call c a stationary point
- singular points: If $c \in [a, b]$ where f' fails to exist, we call c a singular point

Where do extreme values occur?

Let S , the domain of f , contains the point c .

- End points of the interval
- stationary points: If c is a point which $f'(c) = 0$, we call c a stationary point
- singular points: If $c \in [a, b]$ where f' fails to exist, we call c a singular point

Any point of one of these three types is called critical point of f .

What are the extreme values?

Let f be a continuous function defined on an interval $I = [a, b]$.

- Find the critical points of f on I

What are the extreme values?

Let f be a continuous function defined on an interval $I = [a, b]$.

- Find the critical points of f on I
- Evaluate f at each of these critical points. The largest of these values is the maximum value, the smallest is the minimum value.

Monotonicity and Concavity

Let f be defined on an interval I .

- f is increasing on I if for every pair of numbers x_1 and x_2 in I , $x_1 < x_2$ implies $f(x_1) < f(x_2)$.

Monotonicity and Concavity

Let f be defined on an interval I .

- f is increasing on I if for every pair of numbers x_1 and x_2 in I , $x_1 < x_2$ implies $f(x_1) < f(x_2)$.
- f is decreasing on I if for every pair of numbers x_1 and x_2 in I , $x_1 < x_2$ implies $f(x_1) > f(x_2)$.

Monotonicity and Concavity

Let f be defined on an interval I .

- f is increasing on I if for every pair of numbers x_1 and x_2 in I , $x_1 < x_2$ implies $f(x_1) < f(x_2)$.
- f is decreasing on I if for every pair of numbers x_1 and x_2 in I , $x_1 < x_2$ implies $f(x_1) > f(x_2)$.
- f is strictly monotone on I if it is either increasing or decreasing on I

Monotonicity Theorem

Let f be a continuous function on an interval I and differentiable at every interior point of I .

Monotonicity Theorem

Let f be a continuous function on an interval I and differentiable at every interior point of I .

- If $f'(x) > 0$ for all x interior point of I then f is increasing on I .

Monotonicity Theorem

Let f be a continuous function on an interval I and differentiable at every interior point of I .

- If $f'(x) > 0$ for all x interior point of I then f is increasing on I .
- If $f'(x) < 0$ for all x interior point of I then f is decreasing on I .

Concave up and Concave down

Let f be differentiable on an open interval I . We say that f is concave up on I if f' is increasing on I , and we say that f is concave down on I if f' is decreasing on I .

Concave up and Concave down

Let f be differentiable on an open interval I . We say that f is concave up on I if f' is increasing on I , and we say that f is concave down on I if f' is decreasing on I . We call $(c, f(c))$ an inflection point of the graph of f , if f is concave up on one side of c and concave down on the other side.

Concavity Theorem

Let f be twice differentiable on the open interval I .

Concavity Theorem

Let f be twice differentiable on the open interval I .

- If $f''(x) > 0$ for all $x \in I$ then f is concave up on I .

Concavity Theorem

Let f be twice differentiable on the open interval I .

- If $f''(x) > 0$ for all $x \in I$ then f is concave up on I .
- If $f''(x) < 0$ for all $x \in I$ then f is concave down on I .

Local extrema

Let S , the domain of f , contain a certain point c .

Local extrema

Let S , the domain of f , contain a certain point c .

- $f(c)$ is a local maximum value of f on S if there is an interval (a, b) containing c such that $f(c)$ is the maximum value on $S \cap (a, b)$.

Let S , the domain of f , contain a certain point c .

- $f(c)$ is a local maximum value of f on S if there is an interval (a, b) containing c such that $f(c)$ is the maximum value on $S \cap (a, b)$.
- $f(c)$ is a local minimum value of f on S if there is an interval (a, b) containing c such that $f(c)$ is the minimum value on $S \cap (a, b)$.

Let S , the domain of f , contain a certain point c .

- $f(c)$ is a local maximum value of f on S if there is an interval (a, b) containing c such that $f(c)$ is the maximum value on $S \cap (a, b)$.
- $f(c)$ is a local minimum value of f on S if there is an interval (a, b) containing c such that $f(c)$ is the minimum value on $S \cap (a, b)$.
- $f(c)$ is a local extreme value of f if it is either a local maximum value or a local minimum value.

First Derivative Theorem

Let f be a continuous function on an open interval $I = (a, b)$ that contains a critical point c .

First Derivative Theorem

Let f be a continuous function on an open interval $I = (a, b)$ that contains a critical point c .

- If $f'(x) > 0$ for all x in (a, c) and $f'(x) < 0$ for all x in (c, b) then $f(c)$ is a local maximum.

First Derivative Theorem

Let f be a continuous function on an open interval $I = (a, b)$ that contains a critical point c .

- If $f'(x) > 0$ for all x in (a, c) and $f'(x) < 0$ for all x in (c, b) then $f(c)$ is a local maximum.
- If $f'(x) < 0$ for all x in (a, c) and $f'(x) > 0$ for all x in (c, b) then $f(c)$ is a local minimum.

Second Derivative Theorem

Let f' and f'' exist at every point in an open interval $I = (a, b)$ containing c and suppose $f'(c) = 0$.

Second Derivative Theorem

Let f' and f'' exist at every point in an open interval $I = (a, b)$ containing c and suppose $f'(c) = 0$.

- If $f''(c) < 0$ then $f(c)$ is a local maximum of f .

Second Derivative Theorem

Let f' and f'' exist at every point in an open interval $I = (a, b)$ containing c and suppose $f'(c) = 0$.

- If $f''(c) < 0$ then $f(c)$ is a local maximum of f .
- If $f''(c) > 0$ then $f(c)$ is a local minimum of f .